arXiv:math/0403315v3 [math.QA] 18 May 2004

CHARACTER AND DIMENSION FORMULAE FOR GENERAL LINEAR SUPERALGEBRA

YUCAI SU AND R. B. ZHANG

Abstract. The generalized Kazhdan-Lusztig polynomials for the finite dimensional irreducible representations of the general linear superalgebra are computed explicitly. Using the result we establish a one to one correspondence between the set of composition factors of an arbitrary r-fold atypical $\mathfrak{gl}_{m|n}$-Kac-module and the set of composition factors of some r-fold atypical $\mathfrak{gl}_{r|r}$-Kac-module. The result of Kazhdan-Lusztig polynomials is also applied to prove a conjectural character formula put forward by van der Jeugt et al in the late 80s. We simplify this character formula to cast it into the Kac-Weyl form, and derive from it a closed formula for the dimension of any finite dimensional irreducible representation of the general linear superalgebra.

1. Introduction

Formal characters of finite dimensional irreducible representations of complex simple Lie superalgebras encapsulate rich information on the structure of the representations themselves. In his foundational papers [6, 7, 8, 9] on Lie superalgebras, Kac raised the problem of determining the formal characters of finite dimensional irreducible representations of Lie superalgebras, and developed a character formula for the so-called typical irreducible representations. However, the problem turned out to be quite hard for the so-called atypical irreducible representations. In the early 80s Bernstein and Leites [1] gave a formula for the general linear superalgebra, which produces the correct formal characters for the singly atypical irreducible representations [15], but fails for the multiply atypical irreducibles (e.g., the trivial representation is multiply atypical). Since then much further research was done on the problem. For the orthosymplectic superalgebra $\mathfrak{osp}_{2|2n}$, van der Jeugt [13] constructed a character formula for all finite dimensional irreducible representations (which are necessarily singly atypical). There were also partial results and conjectures in other cases. Particularly noteworthy is the conjectural character formula for arbitrary finite dimensional irreducible representations of $\mathfrak{gl}_{m|n}$ put forward by van der Jeugt, Hughes, King and Thierry-Mieg [16], which was the result of extensive research carried out by the authors over several years period. Their formula withstood the tests of large scale computer calculations for a wide range of irreducible representations. However, the full problem of determining the formal characters of finite dimensional irreducible representations of Lie superalgebras remained open until 1995 when Serganova [11, 12] used a combination of geometric and algebraic techniques to obtain a general solution.

Serganova's approach was based on ideas from Kazhdan-Lusztig theory. She introduced some generalized Kazhdan-Lusztig polynomials, the values of which at $q = -1$ determine the formal characters of finite dimensional irreducible representations of Lie superalgebras. Serganova's work was further developed in the papers [18, 17, 2, 3, 4]. Particularly important is the work of Brundan, developed a very practicable algorithm for computing the generalized Kazhdan-Lusztig polynomials, by using quantum

---

2000 *Mathematics Subject Classification.* Primary 17B10.

group techniques. This enables him to gain sufficient knowledge on the generalized Kazhdan-Lusztig polynomials to prove the conjecture of [17] on the composition factors of Kac-modules.

In this paper, we shall further investigate Brundan's algorithm and implement it to compute the generalized Kazhdan-Lusztig polynomials for the finite dimensional irreducible representations of the general linear superalgebra. A closed formula is obtained for the generalized Kazhdan-Lusztig polynomials, which is essentially given in terms of the permutation group of the atypical roots (see Theorem 3.24 for details).

The formula for the generalized Kazhdan-Lusztig polynomials is quite explicit and easy to apply. It leads to a relatively explicit character formula for all the finite dimensional irreducible representations (see Theorem 4.1). By analysing this formula we prove that the conjecture of van der Jeugt et al [16] holds true for all finite dimensional irreducible $\mathfrak{gl}_{m|n}$-modules.

A general fact in the context of Lie superalgebras is that the character formula constructed by using Kazhdan-Lusztig theory is always in the form of an infinite sum. This makes the character formula rather unwieldy to use for, e.g., determining dimensions of finite dimensional irreducible representations. Therefore, it is highly desirable to sum up the infinite series to cast the character formula into the Kac-Weyl form. This is done in Theorem 4.9.

Equipped with Theorem 4.9 we are able to work out the dimension of any finite dimensional irreducible representation of $\mathfrak{gl}_{m|n}$, and the result is given by the closed formula of Theorem 4.14. In the special case of singly atypical irreducible representations, our dimension formula reproduces what one obtains from the Bernstein-Leites character formula [14].

In proving Theorem 3.24 on the Kazhdan-Lusztig polynomials we have introduced the notion of heights of a weight with respect to its atypical roots. This notion proves to be extremely useful. All the results in this paper can be presented using this concept. In particular, the Kazhdan-Lusztig polynomial $K_{\lambda,\mu}(q)$ depends only on the heights of λ and μ with respect to their atypical roots. This latter fact enables one to reduce the study of r-fold atypical $\mathfrak{gl}_{m|n}$-Kac-modules to the study of r-fold atypical $\mathfrak{gl}_{r|r}$-Kac-modules. We state this precisely in Theorem 3.29, which establishes a one to one correspondence between the set of composition factors of an arbitrary r-fold atypical $\mathfrak{gl}_{m|n}$-Kac-module and the set of composition factors of some r-fold atypical $\mathfrak{gl}_{r|r}$-Kac-module.

The organization of the paper is as follows. In Section 2 we present some background material on $\mathfrak{gl}_{m|n}$, which will be used throughout the paper. In Section 3 we investigate the generalized Kazhdan-Lusztig polynomials for finite dimensional irreducible $\mathfrak{gl}_{m|n}$-modules. This section contains two main results, Theorem 3.24 and Theorem 3.29. While Theorem 3.24 gives an explicit formula for the Kazhdan-Lusztig polynomials, Theorem 3.29 establishes a one to one correspondence between the set of composition factors of an arbitrary r-fold atypical Kac-module over $\mathfrak{gl}_{m|n}$ and the set of composition factors of some r-fold atypical Kac-module $\mathfrak{gl}_{r|r}$. In Section 4 we first use Theorem 3.24 to prove the conjectural character formula of van der Jeugt et al [16] (see Theorem 4.2), then we re-write the formula into the Kac-Weyl form (see Theorem 4.9). Finally we derive from Theorem 4.9 a closed formula for the dimension of any finite dimensional irreducible representation of the general linear superalgebra (see Theorem 4.14).

2. Preliminaries

We explain some basic notions of Lie superalgebras here and refer to [7, 5, 16] for more details. We shall work over the complex number field $\mathbb{C}$ throughout the paper. Given a $\mathbb{Z}_2$-graded vector space $W = W_{\bar{0}} \oplus W_{\bar{1}}$, we call $W_{\bar{0}}$ and $W_{\bar{1}}$ the even and odd subspaces, respectively. Define a map $[\ ] : W_{\bar{0}} \cup W_{\bar{1}} \to \mathbb{Z}_2$ by $[w] = \bar{i}$ if $w \in W_{\bar{i}}$. For any two $\mathbb{Z}_2$-graded vector spaces V and W, the space of morphisms $\mathrm{Hom}_{\mathbb{C}}(V, W)$ (in the category of $\mathbb{Z}_2$-graded vector spaces) is also $\mathbb{Z}_2$-graded with $\mathrm{Hom}_{\mathbb{C}}(V, W)_{\bar{k}} = \sum_{\bar{i}+\bar{j}=\bar{k}} \mathrm{Hom}_{\mathbb{C}}(V_{\bar{i}}, W_{\bar{j}})$. We write $\mathrm{End}_{\mathbb{C}}(V)$ for $\mathrm{Hom}_{\mathbb{C}}(V, V)$.

Let $\mathbb{C}^{m|n}$ be the $\mathbb{Z}_2$-graded vector space with even subspace $\mathbb{C}^m$ and odd subspace $\mathbb{C}^n$. Then $\mathrm{End}_{\mathbb{C}}(\mathbb{C}^{m|n})$ with the $\mathbb{Z}_2$-graded commutator forms the *general linear superalgebra*. To describe its structure, we choose a homogeneous basis $\{v_a \,|\, a \in \boldsymbol{I}\}$, for $\mathbb{C}^{m|n}$, where $\boldsymbol{I} = \{1, 2, \ldots, m+n\}$, and v_a is even if $a \le m$, and odd otherwise. The general linear superalgebra relative to this basis of $\mathbb{C}^{m|n}$ will be denoted by $\mathfrak{gl}_{m|n}$, which shall be further simplified to $\mathfrak{g}$ throughout the paper. Let E_{ab} be the matrix unit, namely, the $(m+n) \times (m+n)$-matrix with all entries being zero except that at the (a, b) position which is 1. Then $\{E_{ab} \,|\, a, b \in \boldsymbol{I}\}$ forms a homogeneous basis of $\mathfrak{g}$, with E_{ab} being even if $a, b \le m$, or $a, b > m$, and odd otherwise. For convenience, we define the map $[\ ] : \boldsymbol{I} \to \mathbb{Z}_2$ by $[a] = \begin{cases} \bar{0}, & \text{if } a \le m, \\ \bar{1}, & \text{if } a > m. \end{cases}$ Then the commutation relations of the Lie superalgebra can be written as

$$[E_{ab},\ E_{cd}] \;=\; E_{ad}\delta_{bc} - (-1)^{([a]-[b])([c]-[d])} E_{cb}\delta_{ad}.$$

The upper triangular matrices form a *Borel subalgebra* $\mathfrak{b}$ of $\mathfrak{g}$, which contains the *Cartan subalgebra* $\mathfrak{h}$ of diagonal matrices. Let $\{\epsilon_a \,|\, a \in \boldsymbol{I}\}$ be the basis of $\mathfrak{h}^*$ such that $\epsilon_a(E_{bb}) = \delta_{ab}$. The supertrace induces a *bilinear form* $(\ ,\ ) : \mathfrak{h}^* \times \mathfrak{h}^* \to \mathbb{C}$ on $\mathfrak{h}^*$ such that

$$(\epsilon_a, \epsilon_b) = (-1)^{[a]}\delta_{ab}.$$

Relative to the Borel subalgebra $\mathfrak{b}$, the roots of $\mathfrak{g}$ can be expressed as $\epsilon_a - \epsilon_b,\ a \ne b$, where $\epsilon_a - \epsilon_b$ is even if $[a] + [b] = \bar{0}$ and odd otherwise. The set of the *positive roots* is $\Delta^+ = \{\epsilon_a - \epsilon_b \,|\, a < b\}$, and the set of *simple roots* is $\{\epsilon_a - \epsilon_{a+1} \,|\, a < m+n\}$.

We denote $\boldsymbol{I}^1 = \{1, 2, ..., m\}$ and $\boldsymbol{I}^2 = \{1, 2, ..., n\}$. We also set $\delta_\zeta = \epsilon_{\dot{\zeta}}$ for $\zeta \in \boldsymbol{I}^2$, where we use the notation

$$\dot{\zeta} = \zeta + m.$$

Then the sets of *positive even roots* and *odd roots* are respectively

$$\begin{aligned} \Delta_0^+ &= \{\epsilon_i - \epsilon_j,\, \delta_\zeta - \delta_\eta \,|\, 1 \le i < j \le m,\, 1 \le \zeta < \eta \le n\}, \\ \Delta_1^+ &= \{\epsilon_i - \delta_\zeta \,|\, i \in \boldsymbol{I}^1,\, \zeta \in \boldsymbol{I}^2\}. \end{aligned}$$

The Lie algebra $\mathfrak{g}$ admits a $\mathbb{Z}_2$-consistent $\mathbb{Z}$-grading

$$\mathfrak{g} = \mathfrak{g}_{-1} \oplus \mathfrak{g}_0 \oplus \mathfrak{g}_{+1}, \text{ where } \mathfrak{g}_0 = \mathfrak{g}_{\bar{0}} \cong \mathfrak{gl}(m) \oplus \mathfrak{gl}(n) \text{ and } \mathfrak{g}_{\pm 1} \subset \mathfrak{g}_{\bar{1}},$$

with $\mathfrak{g}_{+1}$ (resp. $\mathfrak{g}_{-1}$) being the nilpotent subalgebra spanned by the odd positive (resp. negative) root spaces. We define a total order on Δ_1^+ by

$$\epsilon_i - \delta_\zeta < \epsilon_j - \delta_\eta \quad \Longleftrightarrow \quad \zeta - i < \eta - j \text{ or } \zeta - i = \eta - j \text{ but } i > j. \tag{2.1}$$

An element in $\mathfrak{h}^*$ is called a *weight*. A weight $\lambda \in \mathfrak{h}^*$ will be written in terms of the $\epsilon\delta$-basis as

$$\lambda = (\lambda_1, ..., \lambda_m \,|\, \lambda_{\dot{1}}, ..., \lambda_{\dot{n}}) = \sum_{i \in \boldsymbol{I}^1} \lambda_i \epsilon_i - \sum_{\zeta \in \boldsymbol{I}^2} \lambda_{\dot{\zeta}} \delta_\zeta, \tag{2.2}$$

where we have adopted an unusual (but convenient) sign convention for λ_i's. Thus $\lambda_i=(\lambda,\epsilon_i)$, called the *i-th entry of* λ for $i\in\boldsymbol{I}^1$, and $\lambda_{\dot\zeta}=(\lambda,\delta_\zeta)$, called the $\dot\zeta$*-th entry of* λ for $\zeta\in\boldsymbol{I}^2$. A weight λ is called

$$\begin{aligned}
&\textit{integral} &&\Longleftrightarrow\quad \lambda_i,\lambda_{\dot\zeta}\in\mathbb{Z}\quad\text{for}\ \ i\in\boldsymbol{I}^1,\ \zeta\in\boldsymbol{I}^2; &&(2.3)\\
&\textit{dominant} &&\Longleftrightarrow\quad 2(\Lambda,\,\alpha)/(\alpha,\,\alpha)\geq 0\quad\text{for all positive even roots } \alpha \text{ of } \mathfrak{g}\text{, namely,}\\
& && \qquad\ \ \lambda_1\geq ...\geq\lambda_m,\qquad \lambda_{\dot 1}\leq ...\leq\lambda_{\dot n}. &&(2.4)
\end{aligned}$$

Denote by P (resp. P_+) the set of integral (resp. dominant integral) weights. Using notation (2.2), P coincides with the set of the $m|n$-tuples of integers, thus P is also denoted by $\mathbb{Z}^{m|n}$, and P_+ by $\mathbb{Z}^{m|n}_+$.

Let ρ_0 (resp. ρ_1) be half the sum of positive even (resp. odd) roots, and let $\rho=\rho_0-\rho_1$. Then

$$\begin{aligned}
\rho_0&=\frac{1}{2}\Big(\sum_{i=1}^{m}(m-2i+1)\epsilon_i+\sum_{\zeta=1}^{n}(n-2\zeta+1)\delta_\zeta\Big)\\
&=\frac{1}{2}(m-1,m-3,...,1-m\,|\,1-n,3-n,...,n-1), \qquad (2.5)\\
\rho_1&=\frac{1}{2}\Big(n\sum_{i=1}^{m}\epsilon_i-m\sum_{\zeta=1}^{n}\delta_\zeta\Big)=\frac{1}{2}(n,...,n\,|\,m,...,m),\\
\rho&=\rho'-\tfrac{m+n+1}{2}\mathbf{1},\ \text{ where } \rho'=(m,...,2,1\,|\,1,2,...,n),\ \mathbf{1}=(1,...,1\,|\,1,...,1).
\end{aligned}$$

For all purposes, we can replace ρ by ρ'. Therefore, from here on we shall denote

$$\rho=(m,...,2,1\,|\,1,2,...,n). \tag{2.6}$$

Let $W=\mathit{Sym}_m\times\mathit{Sym}_n$ be the *Weyl group* of $\mathfrak{g}$, where Sym_m is the *symmetric group of degree m*. We define the *dot action* of W on P by

$$w\cdot\mu=w(\mu+\rho)-\rho\qquad\text{for}\quad w\in W,\ \mu\in P. \tag{2.7}$$

An integral weight λ is called

- *regular* or *non-vanishing* (in sense of [5, 16]) if it is W-conjugate under the dot action to a dominant weight (which is denoted by λ^+ throughout the paper);
- *vanishing* otherwise (since the right-hand side of (2.13) is vanishing in this case, cf. (4.8)).

Obviously,

$$\lambda\ \text{ is regular }\ \Longleftrightarrow\ \begin{cases}\lambda_1+m,\ \lambda_2+m-1,\ ...,\ \lambda_m+1 & \text{are all distinct, and}\\ \lambda_{\dot 1}+1,\ \lambda_{\dot 2}+2,\ ...,\ \lambda_{\dot n}+n & \text{are all distinct.}\end{cases} \tag{2.8}$$

Let λ in (2.2) be a regular weight. A positive odd root $\epsilon_i-\delta_\zeta$ is an *atypical root of* λ if

$$(\lambda+\rho,\epsilon_i-\delta_\zeta)=(\lambda_i+m+1-i)-(\lambda_{\dot\zeta}+\zeta)=0. \tag{2.9}$$

Denote by Γ_λ the set of atypical roots of λ (cf. (3.1) and (3.2)):

$$\Gamma_\lambda=\{\epsilon_i-\delta_\zeta\,|\,(\lambda+\rho,\epsilon_i-\delta_\zeta)=0\}. \tag{2.10}$$

Set $r=\#\Gamma_\lambda$. We also denote $\#\lambda=r$, called the *degree of atypicality of* λ. A weight λ is called

- *typical* if $r=0$;
- *atypical* if $r>0$ (in this case λ is also called an *r-fold atypical weight*).

Let $V=\oplus_{\mu\in\mathfrak{h}^*}V_\mu$ be a *weight module* over $\mathfrak{g}$, where

$$V_\mu=\{v\in V\,|\,hv=\mu(h)v,\,\forall\,h\in\mathfrak{h}\}\quad\text{with}\quad\dim V_\mu<\infty,$$

is the weight space of weight μ. The *character* $\mathrm{ch}V$ is defined to be

$$\mathrm{ch}V=\sum_{\mu\in\mathfrak{h}^*}(\dim V_\mu)e^\mu, \tag{2.11}$$

where e^μ is the *formal exponential*, which will be regarded as an element of an additive group isomorphic to $\mathfrak{h}^*$ under $\mu\mapsto e^\mu$. Then $\mathrm{ch}V$ is an element of the *completed group algebra*

$$\varepsilon=\Big\{\sum_{\mu\in\mathfrak{h}^*}a_\mu e^\mu\;\Big|\;a_\mu\in\mathbb{C},\,a_\mu=0\text{ except }\mu\text{ is in a finite union of }\mathcal{Q}_\lambda\Big\}, \tag{2.12}$$

where for $\lambda\in\mathfrak{h}^*$,

$$\mathcal{Q}_\lambda=\Big\{\lambda-\sum_{\alpha\in\Delta^+}i_\alpha\alpha\in\mathfrak{h}^*\;\Big|\;i_\alpha\in\mathbb{Z}_+\Big\}.$$

For every integral dominant weight λ, we denote by $V^{(0)}(\lambda)$ the finite-dimensional irreducible $\mathfrak{g}_0$-module with highest weight λ. Extend it to a $\mathfrak{g}_0\oplus\mathfrak{g}_{+1}$-module by putting $\mathfrak{g}_{+1}V^{(0)}(\lambda)=0$. Then the *Kac-module* $\overline{V}(\lambda)$ is the induced module

$$\overline{V}(\lambda)=\mathrm{Ind}_{\mathfrak{g}_0\oplus\mathfrak{g}_{+1}}^{\mathfrak{g}}V^{(0)}(\lambda)\cong U(\mathfrak{g}_{-1})\otimes_{\mathbb{C}}V^{(0)}(\lambda).$$

Denote by $V(\lambda)$ the *irreducible module* with highest weight λ (which is the unique irreducible quotient module of $\overline{V}(\lambda)$).

The following result is due to Kac [8, 9]:

Theorem 2.1. *If λ is a dominant integral typical weight, then $V(\lambda)=\overline{V}(\lambda)$, and*

$$\mathrm{ch}V(\lambda)=\mathrm{ch}\overline{V}(\lambda)=\frac{L_1}{L_0}\sum_{w\in W}\epsilon(w)e^{w(\lambda+\rho)}, \tag{2.13}$$

where $\epsilon(w)$ is the signature of $w\in W$, and

$$L_0=\prod_{\alpha\in\Delta_0^+}(e^{\alpha/2}-e^{-\alpha/2}),\quad L_1=\prod_{\beta\in\Delta_1^+}(e^{\beta/2}+e^{-\beta/2}). \tag{2.14}$$

Since any finite-dimensional irreducible $\mathfrak{g}$-module is either a typical module or is a tensor module of $V(\lambda)$ with one-dimensional module for some $\lambda\in P_+$, in the rest of the paper there is no loss of generality in restricting our attention to integral weights λ.

3. Kazhdan-Lusztig polynomials

3.1. **The height vectors and the c-relationship.** Let $\lambda\in\mathfrak{h}^*$ be an r-fold atypical dominant integral weight:

$$\lambda=(\lambda_1,...,\overset{\gamma_r}{\lambda_{m_r}},...,\lambda_i,...,\overset{\gamma_1}{\lambda_{m_1}},...,\lambda_m\;\big|\;\lambda_{\dot 1},...,\overset{\gamma_1}{\lambda_{\dot n_1}},...,\lambda_{\dot\zeta},...,\overset{\gamma_r}{\lambda_{\dot n_r}},...,\lambda_{\dot n}), \tag{3.1}$$

with the set of atypical roots (cf. (2.9), (2.10) and (2.1))

$$\Gamma_\lambda=\{\gamma_1,...,\gamma_r\},\qquad\text{where }\;\gamma_1=\epsilon_{m_1}-\delta_{n_1}<...<\gamma_r=\epsilon_{m_r}-\delta_{n_r}, \tag{3.2}$$

and $m_r<...<m_1,\;n_1<...<n_r$. We call γ_s the *s-th atypical root of* λ for $s=1,...,r$. For convenience, we introduce the notation λ^ρ for the *ρ-translation of* λ:

$$\lambda^\rho=\lambda+\rho. \tag{3.3}$$

Thus $\{(m_s, n_s) \mid s = 1, ..., r\}$ is the maximal set of pairs (i, ζ) satisfying $\lambda^\rho_i = \lambda^\rho_{\dot\zeta}$ by (2.9). We define the *atypical tuple of* λ

$$\mathrm{aty}_\lambda = (\lambda^\rho_{\dot n_1}, ..., \lambda^\rho_{\dot n_r}) = (\lambda_{\dot n_1} + n_1, ..., \lambda_{\dot n_r} + n_r) \in \mathbb{Z}^r, \tag{3.4}$$

and call the s-th entry of aty_λ the *s-th atypical entry of* λ for $s = 1, ..., r$. We also define the *typical tuple of* λ

$$\mathrm{typ}_\lambda \in \mathbb{Z}^{m-r|n-r} \tag{3.5}$$

to be the element obtained from λ^ρ by deleting all entries $\lambda^\rho_{m_s}$, $\lambda^\rho_{\dot n_s}$ for $s = 1, ..., r$. Thus all entries of typ_λ, called the *typical entries of* λ, are distinct by (2.8).

Definition 3.1. Corresponding to each atypical root γ_s of λ, we define the *γ_s-height of* λ (the *height of* λ *with respect to the s-th atypical root*)

$$h_s(\lambda) = \lambda_{m_s} - n_s + s \quad \text{for} \quad s = 1, ..., r. \tag{3.6}$$

We also introduce the *height vector of* λ and the *height of* λ respectively:

$$h(\lambda) = (h_1(\lambda), ..., h_r(\lambda)), \qquad |h(\lambda)| = \sum_{s=1}^{r} h_s(\lambda). \tag{3.7}$$

Remark 3.2. As we shall see later, the concept of heights of a weight with respect to its atypical roots is extremely useful. In fact, the Kazhdan-Lusztig polynomials are completely determined by the height vectors of the weights involved (see Theorem 3.24 and Theorem 3.29).

Example 3.3. Suppose λ is the weight

$$\lambda = \big(7, \overset{4}{6}, 5, \overset{3}{5}, \overset{2}{3}, 3, 2, \overset{1}{2}, 0 \,\big|\, 1, \overset{1}{2}, 3, \overset{2}{4}, 4, \overset{3}{5}, \overset{4}{7}, 7\big) \in \mathbb{Z}^{9|8}_+, \tag{3.8}$$

where we put a label t over a pair of entries to indicate that they are the entries associated with the t-th atypical root. Then

$$\lambda^\rho = \big(16, \underline{14}, 12, \underline{11}, \underline{8}, 7, 5, \underline{4}, 1 \,\big|\, 2, \underline{4}, 6, \underline{8}, 9, \underline{11}, \underline{14}, 15\big)\,, \tag{3.9}$$

$$\mathrm{aty}_\lambda = (4, 8, 11, 14), \tag{3.10}$$

$$\mathrm{typ}_\lambda = (16, 12, 7, 5, 1 \,|\, 2, 6, 9, 15), \tag{3.11}$$

$$h(\lambda) = (1, 1, 2, 3), \tag{3.12}$$

where the underlined integers in λ^ρ are entries of atypical tuple aty_λ.

We define the partial order '$\preccurlyeq$' on P_+ for $\lambda, \mu \in P_+$ by

$$\mu \preccurlyeq \lambda \quad \Longleftrightarrow \quad \#\mu = \#\lambda,\ \ \mathrm{aty}_\mu \le \mathrm{aty}_\lambda \ \text{ and } \ \mathrm{typ}_\mu = \mathrm{typ}_\lambda, \tag{3.13}$$

where the partial order "$\le$" on $\mathbb{Z}^r$ is defined for $a = (a_1, ..., a_r)$, $b = (b_1, ..., b_r) \in \mathbb{Z}^r$ by

$$a \le b \quad \Longleftrightarrow \quad a_i \le b_i, \ \ \forall\, i \in [1, r]. \tag{3.14}$$

Now suppose μ is another r-fold atypical dominant integral weight with atypical roots:

$$\gamma'_1 = \epsilon_{m'_1} - \delta_{n'_1} < ... < \gamma'_r = \epsilon_{m'_r} - \delta_{n'_r}. \tag{3.15}$$

We define (cf. (3.22) below)

$$\ell_s(\lambda, \mu) = h_s(\lambda) - h_s(\mu) = \lambda_{m_s} - \mu_{m'_s} + n'_s - n_s \quad \text{ for } \ s \in [1, r], \tag{3.16}$$

and define the *length between* λ *and* μ to be

$$\ell(\lambda, \mu) = \sum_{s=1}^{r} \ell_s(\lambda, \mu) = |h(\lambda)| - |h(\mu)|. \tag{3.17}$$

(In general $\ell(\lambda,\mu)$ is not necessarily non-negative, but when $\mu \preccurlyeq \lambda$, it is indeed non-negative.)

Remark 3.4. *The height* $|h(\lambda)|$ *of* λ *turns out to be the absolute length* $\ell(\lambda)$ *defined by Brundan in* [2, §3-g]*, and the length* $\ell(\lambda,\mu)$ *coincides with the length* $\ell(\mu,\lambda)$ *defined in* [2, §3-g]*.*

For $1 \le s \le t \le r$, we denote

$$d_{s,t}(\lambda) = h_t(\lambda) - h_s(\lambda) = \lambda_{m_t} - \lambda_{m_s} - n_t + n_s + t - s. \tag{3.18}$$

The fact that $d_{s,t}(\lambda)$ is non-negative is not obvious, but one can observe that it is the number of integers between the s-th atypical entry $\lambda^\rho_{\dot n_s}$ and the t-th atypical entry $\lambda^\rho_{\dot n_t}$ which are not entries of λ^ρ, namely (cf. (3.20), (3.21) and (3.22))

$$d_{s,t}(\lambda) = \#\left([\lambda^\rho_{\dot n_s}, \lambda^\rho_{\dot n_t}]\backslash \mathrm{Set}(\lambda^\rho)\right). \tag{3.19}$$

Because of this fact, we call $d_{s,t}(\lambda)$ the *distance between two atypical roots* γ_s *and* γ_t *of* λ. Here and below we use the notation

$$\mathrm{Set}(\mu) = \text{ the set of the entries of a weight } \mu, \tag{3.20}$$

and the notation

$$[i,j] = \begin{cases} \{k \in \mathbb{Z} \,|\, i \le k \le j\} & \text{if } i \le j, \\ \emptyset & \text{otherwise,} \end{cases} \qquad \text{for} \quad i,j \in \mathbb{Z}. \tag{3.21}$$

(There will be no danger of confusing this notation with the Lie bracket as the later will not be used in the remainder of the paper.)

One can generalize (3.19) to obtain the following proposition, which will be used in the proof of Theorem 3.29.

Proposition 3.5. *Let* λ,μ *be* r*-fold atypical weights with* $\mu \preccurlyeq \lambda$*. For any* $s,t \in [1,r]$ *with* $\mu^\rho_{\dot n'_s} \le \lambda^\rho_{\dot n_t}$*, we have*

$$\#\left([\mu^\rho_{\dot n'_s}+1, \lambda^\rho_{\dot n_t}]\backslash \mathrm{Set}(\mathrm{typ}_\lambda)\right) = h_t(\lambda) - h_s(\mu) + t - s. \tag{3.22}$$

Proof. By (3.6) and the fact that $\lambda^\rho_{m_s} = \lambda^\rho_{\dot n_s}$, the right-hand side of (3.22) is equal to

$$\lambda_{\dot n_t} + m_t - (\mu_{\dot n'_s} + m'_s) + 2(t-s) = \lambda^\rho_{\dot n_t} - \mu^\rho_{\dot n'_s} - (m'_s - m_t + n_t - n'_s) - 2(s-t).$$

Thus (3.22) is equivalent to

$$\#\left([\mu^\rho_{\dot n'_s}+1, \lambda^\rho_{\dot n_t}] \cap \mathrm{Set}(\mathrm{typ}_\lambda)\right) = m'_s - m_t + n_t - n'_s + 2(s-t). \tag{3.23}$$

Note from (3.13) that $\mathrm{typ}_\lambda = \mathrm{typ}_\mu$. By (2.4), (2.8) and (3.1), in λ^ρ (resp., μ^ρ), the number of typical entries to the left of entry $\lambda^\rho_{\dot m_t}$ (resp., $\mu^\rho_{m'_s}$) is $m_t - 1 - r + t$ (resp., $m'_s - 1 - r + s$). Thus the number of typical entries to the left of the entry $\mu^\rho_{m'_s}$ which are in $[\mu^\rho_{m'_s}+1, \lambda^\rho_{m_t}]$ is $m'_s - m_t + s - t$. Similarly, the number of typical entries to the right of the entry $\mu^\rho_{m'_s}$ which are in $[\mu^\rho_{\dot n'_s}+1, \lambda^\rho_{\dot n_t}]$ is $n_t - n'_s + s - t$. Hence we obtain (3.23). □

For any two distinct atypical roots γ_s and γ_t, there exist unique positive even roots α_{st} and β_{st} such that $(\alpha_{st},\beta_{st}) = 0$, and the composite of the actions of the Weyl group elements $\sigma_{\alpha_{st}}$, $\sigma_{\beta_{st}}$, respectively corresponding to reflections with respect to these even roots, send γ_s to γ_t, namely, $\gamma_t = \sigma_{\alpha_{st}}\sigma_{\beta_{st}}(\gamma_s)$. In terms of explicit formulae, we have

$$\alpha_{st} = \epsilon_{m_t} - \epsilon_{m_s}, \quad \beta_{st} = \delta_{n_s} - \delta_{n_t} \in \Delta_0^+ \quad \text{for} \quad 1 \le s < t \le r. \tag{3.24}$$

The concept of c-relationship defined below was first introduced in [5] from a different point of view.

Definition 3.6. For $s \leq t$, two atypical roots γ_s, γ_t of λ are called *c-related* (in the sense of [5]) or *connected* (in the sense of [17]) if $s = t$ or $d_{s,t}(\lambda) < t - s$. The later relation has the nice interpretation that

> *the distance (i.e., $d_{s,t}(\lambda)$) between two atypical roots is smaller than the number (i.e., $t - s$) of atypical roots between them.*

Relation $d_{s,t}(\lambda) < t - s$ is also equivalent to

$$\lambda_{m_t} - \lambda_{m_s} < n_t - n_s, \tag{3.25}$$

or in terms of weights,

$$\frac{2(\lambda, \alpha_{st})}{(\alpha_{st}, \alpha_{st})} + \frac{2(\lambda, \beta_{st})}{(\beta_{st}, \beta_{st})} < \frac{2(\rho, \alpha_{st})}{(\alpha_{st}, \alpha_{st})} + \frac{2(\rho, \beta_{st})}{(\beta_{st}, \beta_{st})}. \tag*{(3.25)$'$}$$

We define

$$c_{s,t}(\lambda) = \begin{cases} 1 & \text{if the atypical roots } \gamma_s, \gamma_t \text{ of } \lambda \text{ are } c\text{-related,} \\ 0 & \text{otherwise.} \end{cases} \tag{3.26}$$

We also define

$$\widehat{c}_{s,t}(\lambda) = \begin{cases} 1 & \text{if } \ c_{s,s}(\lambda) = c_{s,s+1}(\lambda) = ... = c_{s,t}(\lambda) = 1, \\ 0 & \text{otherwise,} \end{cases} \tag{3.27}$$

and we say that γ_s, γ_t *are strongly c-related* if $\widehat{c}_{s,t}(\lambda) = 1$.

Note that c-relationship is reflexive and transitive but not symmetric ($c_{t,s}(\lambda)$ is not defined when $t > s$).

Example 3.7. Let λ be the weight in (3.8). By (3.25),

$c_{1,2}(\lambda) = 1$ since $\lambda_{m_2} - \lambda_{m_1} = 3 - 2 < 4 - 3 = n_2 - n_1$;
$c_{1,3}(\lambda) = 1$ since $\lambda_{m_3} - \lambda_{m_1} = 5 - 2 < 6 - 2 = n_3 - n_1$;
$c_{1,4}(\lambda) = 1$ since $\lambda_{m_4} - \lambda_{m_1} = 6 - 2 < 7 - 2 = n_4 - n_1$; and
$c_{s,t}(\lambda) = 0$ for any other pair (s, t).

Thus

$$\widehat{c}_{1,2}(\lambda) = \widehat{c}_{1,3}(\lambda) = \widehat{c}_{1,4}(\lambda) = 1 \text{ and } \widehat{c}_{s,t}(\lambda) = 0 \text{ for any other pair } (s, t).$$

Remark 3.8. If λ is regular but not necessarily dominant, we shall generalize the notions $c, \widehat{c}, d, h$ to λ by defining them with respect to the dominant weight λ^+. For instance, $c_{s,t}(\lambda) = c_{s,t}(\lambda^+)$. Sometimes even if λ is not regular but lexical (in the sense of Definition 3.9), one can still define the c-relationship by using the distance (3.18).

3.2. **Lexical weights.** Let λ be an r-fold atypical regular weight (not necessarily dominant) with the set $\Gamma_\lambda = \{\gamma_1, ..., \gamma_r\}$ of atypical roots ordered according to (3.2). We call λ *lexical* if its atypical tuple aty_λ is lexical in the following sense:

Definition 3.9. An element $a = (a_1, ..., a_r) \in \mathbb{Z}^r$ is called *lexical* if

$$a_1 \leq ... \leq a_r. \tag{3.28}$$

The two sets P_r and P_r^{Lex} to be defined below will be frequently used throughout this section.

Definition 3.10. We denote by P_r the set of r-fold atypical regular weights λ of the form (3.1) such that the atypical roots of λ can be ordered as in (3.2) and that the typical tuple $\mathrm{typ}_\lambda \in \mathbb{Z}_+^{m-r|n-r}$ is dominant as a weight for $\mathfrak{gl}_{m-r|n-r}$.

We denote by P_r^{Lex} the subset of P_r consisting of the lexical weights of P_r.

3.3. **The r-tuple of positive integers associated with λ.** The r-tuple associated with $\lambda \in P_r^{\mathrm{Lex}}$ defined below was first introduced in [17].

Definition 3.11. Define the r-tuple $(k_1, ..., k_r)$ of positive integers associated with $\lambda \in P_r^{\mathrm{Lex}}$ in the following way: each k_s is the smallest positive integer such that

$$(\lambda + \theta_t k_t \gamma_t) + k_s\gamma_s \quad \text{is regular for all } \ t = s+1, s+2, ..., r \text{ and } \theta_t \in \{0,1\}. \tag{3.29}$$

The following lemma gives a way to compute k_s's. For $\lambda \in P_r^{\mathrm{Lex}}$ and $s \in [1, r]$, we set

$$\max{}_s^\lambda = \max\{p \in [s,r] \mid \widehat{c}_{s,p}(\lambda) = 1\} \tag{3.30}$$

to be the maximal number $p \in [s, r]$ satisfying the condition that the p-th atypical root γ_p of λ is strongly c-related to γ_s. One immediately sees that

$$\max{}_t^\lambda \le \max{}_s^\lambda \quad \text{ for any } t \text{ with } s \le t \le \max{}_s^\lambda. \tag{3.31}$$

Lemma 3.12. *Let $\lambda \in P_r^{\mathrm{Lex}}$.*

1. *For $s \in [1,r]$, k_s is the integer such that $\lambda^\rho_{\dot n_s}+k$ is the $(\max_s^\lambda + 1 - s)$-th smallest integer bigger than $\lambda^\rho_{\dot n_s}$ and not in the entry set $\mathrm{Set}(\lambda^\rho)$, i.e.,*

$$k_s = \min\left\{ k > 0 \;\middle|\; \#\left([\lambda^\rho_{\dot n_s}, \lambda^\rho_{\dot n_s} + k]\backslash \mathrm{Set}(\lambda^\rho)\right) = \max{}_s^\lambda + 1 - s \right\}. \tag{3.32}$$

2. *The tuple $(k_r, ..., k_1)$ is the lexicographically smallest tuple of positive integers such that for all $\theta = (\theta_1, ..., \theta_r) \in \{0,1\}^r$, $\lambda + \sum_{s=1}^r \theta_s k_s \gamma_s$ is regular. Thus $(k_r, ..., k_1)$ is the tuple satisfying* [2, Main Theorem].

Proof. (1) Denote by k'_s the right-hand side of (3.32). Obviously, k'_r is the smallest positive integer such that $\lambda + k'_r\gamma_r$ is regular because for any $0 < k < k'_r$, by definition the integer $\lambda^\rho_{\dot n_r} + k$ which is equal to $\lambda^\rho_{m_r} + k$ already appears in the entry set $\mathrm{Set}(\lambda^\rho)$ and thus $\lambda + k'_r\gamma_r$ is not regular by (2.8).

If $s < r$, by induction on $\max_s^\lambda$, it is straightforward to see that $\lambda^\rho_{\dot n_s} + k'_s$ is the smallest integer (bigger than $\lambda^\rho_{\dot n_s}$) which is not in $\mathrm{Set}(\lambda^\rho + \theta_t k'_t \gamma_t)$ for $\theta_t \in \{0,1\}$ and $s < t \le r$. Thus k'_s is the smallest positive integer satisfying (3.29).

(2) Similarly, $\lambda^\rho_{\dot n_s} + k'_s$ is also the smallest integer (bigger than $\lambda^\rho_{\dot n_s}$) which is not in $\mathrm{Set}(\lambda^\rho + \sum_{t=s}^r \theta_t k'_t\gamma_t)$ for $\theta = (\theta_s, ..., \theta_r) \in \{0,1\}^{r+1-s}$, i.e, $(k'_r, ..., k'_s)$ is the lexicographically smallest tuple of positive integers such that for all $\theta = (\theta_s, ..., \theta_r) \in \{0,1\}^{r+1-s}$, the weight $\lambda + \sum_{t=s}^r \theta_t k'_t \gamma_t$ is regular. □

Lemma 3.12(1) allows us to compute k_s by the following procedure.

Procedure 3.13. *First set $S = \mathrm{Set}(\lambda^\rho)$. Suppose we have computed $k_r, ..., k_{s+1}$. To compute k_s, we count the numbers in the set S starting with $\lambda^\rho_{\dot n_s}$ until we find a number, say k, not in S. Then $k_s = k - \lambda^\rho_{\dot n_s}$. Now add k into the set S, and continue.*

Example 3.14. Let λ be given in (3.8). Using the above procedure we obtain $(k_4, k_3, k_2, k_1) = (3, 2, 2, 14)$ (cf. (3.9)).

Remark 3.15. If $\lambda \in P_r$ (not necessarily in P_r^{Lex}), we can still compute k_s by the above procedure, but the difference lies in that the k_s's are computed not in the order $s = r, ..., 1$, but in the order that each time we compute k_s with $\lambda^\rho_{\dot n_s}$ being the largest among all those $\lambda^\rho_{\dot n_s}$'s, the corresponding k_s's of which are not yet computed.

3.4. **Raising operators.** Following [2], we define the *raising operator* $R_{m_s,\dot{n}_s}$ on P_r by

$$R_{m_s,\dot{n}_s}(\lambda) = \lambda + k_s\gamma_s \quad \text{for} \ \ \lambda \in P_r \ \ \text{and} \ \ s \in [1,r], \tag{3.33}$$

where γ_s, k_s are defined in (3.2) and Definition 3.11 respectively. Obviously, for $\lambda, \mu \in P_r^{\text{Lex}}$,

$$\mu = R_{m_s,\dot{n}_s}(\lambda) \implies \text{typ}_\lambda = \text{typ}_\mu \quad (\text{cf. } (3.5)). \tag{3.34}$$

Denote $\mathbb{N} = \{0, 1, ...\}$, and let $\theta = (\theta_1, ..., \theta_r) \in \mathbb{N}^r$. We define

$$R'_\theta(\lambda) = (R^{\theta_1}_{m_1,\dot{n}_1} \circ \cdots \circ R^{\theta_r}_{m_r,\dot{n}_r}(\lambda))^+, \tag{3.35}$$

where in general μ^+ denotes the unique dominant element which is W-conjugate under the dot action to μ (cf. (2.7)).

Let $\mu \in P_r$ be another weight with atypical roots $\gamma'_s = \epsilon_{m'_s} - \delta_{n'_s}$, $1 \le s \le r$, being as in (3.15). Then we have (cf. [2, §3-f])

$$\mu \preccurlyeq \lambda \iff \#\mu = \#\lambda =: r, \quad \text{and} \ \ \exists\, \theta \in \mathbb{N}^r \ \ \text{with} \ \ R'_\theta(\mu) = \lambda. \tag{3.36}$$

For convenience, we denote

$$\mu \prec\!\!\prec \lambda \quad \text{if} \ \ \text{typ}_\mu = \text{typ}_\lambda \ \ \text{and} \ \ \max\{\mu^\rho_{\dot{n}'_s} \,|\, s \in [1,r]\} \le \min\{\lambda^\rho_{\dot{n}_s} \,|\, s \in [1,r]\}. \tag{3.37}$$

3.5. **Definitions of S^λ and $S^{\lambda,\mu}$.** The symmetric group Sym_r of degree r acts on $\mathbb{Z}^r$ by permuting entries. This action induces an action on P_r given by

$$\sigma(\lambda) = (\lambda_1, ..., \underbrace{\lambda_{m_{\sigma(r)}}, ..., \lambda_i, ..., \lambda_{m_{\sigma(1)}}}_{\text{atypical entries permuted}}, ..., \lambda_m \,|\, \lambda_{\dot{1}}, ..., \underbrace{\lambda_{\dot{n}_{\sigma(1)}}, ..., \lambda_{\dot{\zeta}}, ..., \lambda_{\dot{n}_{\sigma(r)}}}_{\text{atypical entries permuted}}, ..., \lambda_{\dot{n}}), \tag{3.38}$$

for $\sigma \in Sym_r$ and $\lambda \in P_r$. With this action on P_r, the group Sym_r can be regarded as a subgroup of W, such that every element is of even parity. Thus we also have the *dot action*

$$\sigma \cdot \lambda = \sigma(\lambda + \rho) - \rho \quad \text{for} \ \ \sigma \in Sym_r. \tag{3.39}$$

Definition 3.16. Let $\lambda, \mu \in P_r^{\text{Lex}}$. Define S^λ to be the subset of the symmetric group Sym_r consisting of permutations σ which do not change the order of $s < t$ when the atypical roots γ_s and γ_t of λ are strongly c-related. That is,

$$S^\lambda = \{\sigma \in Sym_r \,|\, \sigma^{-1}(s) < \sigma^{-1}(t) \ \ \text{for all} \ \ s < t \ \ \text{with} \ \ \widehat{c}_{s,t}(\lambda) = 1\}, \tag{3.40}$$

where $\widehat{c}_{s,t}(\lambda)$ is defined in (3.27). We also define $S^{\lambda,\mu}$ to be subset of S^λ consisting of permutations σ such that $\mu \preccurlyeq \sigma \cdot \lambda$, namely

$$S^{\lambda,\mu} = \{\sigma \in S^\lambda \mid \mu \preccurlyeq \sigma \cdot \lambda\}. \tag{3.41}$$

For convenience, we denote

$$\widetilde{S}^{\lambda,\mu} = \{\sigma \in Sym_r \mid \mu \preccurlyeq \sigma \cdot \lambda\}. \tag{3.42}$$

Thus $S^{\lambda,\mu} = S^\lambda \cap \widetilde{S}^{\lambda,\mu}$.

Example 3.17. If λ is the weight in (3.8), then $S^\lambda = \{\sigma \in Sym_4 \,|\, \sigma(1) = 1\} \cong Sym_3$, is a subgroup of Sym_4 (however in general S^λ is not a subgroup).

Let $\ell(\sigma)$ denote the normal *length function on Sym_r*, namely

$$\ell(\sigma) = \sum_{s=1}^{r} \ell(\sigma, s), \quad \text{where} \tag{3.43}$$

$$\ell(\sigma, s) = \#\{t > s \,|\, \sigma(t) < \sigma(s)\} \quad \text{for} \quad s = 1, ..., r. \tag{3.44}$$

For any subset $B \subset Sym_r$, we define the *q-length function of B* by:

$$B(q) = \sum_{\sigma \in B} q^{\ell(\sigma)}. \tag{3.45}$$

Proposition 3.18. *Let $\lambda, \mu \in P_r^{\mathrm{Lex}}$ with $\mu \preccurlyeq \lambda$. We have*

$$\begin{aligned} Sym_r(q) &= (1+q)(1+q+q^2)\cdots(1+q+\cdots+q^{r-1}) = \prod_{s=1}^{r} \frac{q^s-1}{q-1}, && (3.46) \\ S^{\lambda}(q) &= \prod_{s=1}^{r}(q^s-1)\Big/\prod_{s=1}^{r}(q^{\max_s^{\lambda}-s+1}-1), && (3.47) \\ \widetilde{S}^{\lambda,\mu}(q) &= (1+q+\cdots+q^{r-i_r})(1+q+\cdots+q^{r-1-i_{r-1}})\cdots \\ &= \prod_{s=1}^{r} \frac{q^{s+1-i_s}-1}{q-1}, && (3.48) \end{aligned}$$

where

$$i_s = \min\{i \in [1,r] \,|\, \mu^{\rho}_{\dot{n}'_s} \le \lambda^{\rho}_{\dot{n}_i}\} \quad \textit{for} \ \ s \in [1,r]. \tag{3.49}$$

Proof. First we compute $\widetilde{S}^{\lambda,\mu}(q)$. Elements $\sigma \in \widetilde{S}^{\lambda,\mu}$ can be easily described as follows (cf. Example 3.20): for each $s = r, ..., 1$, suppose for all $t > s$, $\sigma(t)$ have been chosen, then $\sigma(s)$ can be any of $s - i_s + 1$ integers $r, r-1, ..., i_s$ which have not yet been taken by the $\sigma(t)$'s for $t > s$. We order the elements of $\{r, ..., i_s\} \backslash \{\sigma(r), ..., \sigma(s+1)\}$ in descending order, and denote by

$$\{r, ..., i_s\} \backslash \{\sigma(r), ..., \sigma(s+1)\} = \{x_1 > ... > x_{s-i_s+1}\}. \tag{3.50}$$

Then each choice of $\sigma(s) = x_k$ for $k = 1, ..., s - i_s + 1$ contributes $k - 1$ to the length $\ell(\sigma)$. Thus we have (3.48). Since $Sym_r = \widetilde{S}^{\lambda,\mu}$ for any $\lambda, \mu \in P_r^{\mathrm{Lex}}$ with $\mu \prec\!\prec \lambda$ (cf. (3.42)), and when $\mu \prec\!\prec \lambda$, all i_s's are equal to 1, we obtain (3.46), which is a well-known formula.

Consider S^{λ} defined in (3.40), which can be re-written as

$$S^{\lambda} = \{\sigma \in Sym_r \,|\, \sigma^{-1}(s) < \sigma^{-1}(t) \text{ for all } s, t \text{ with } s < t \le \max_s^{\lambda}\}.$$

Since for each $s = 1, ..., r$, we cannot change the order of s and t for $s < t \le \max_s^{\lambda}$, we shall remove the factor $1 + q + \cdots + q^{\max_s^{\lambda} - s}$ from $Sym_r(q)$. Thus we obtain (3.47). □

Similar to (3.48), we also have

$$\widetilde{S}^{\lambda,\mu}(q) = \prod_{s=1}^{r} \frac{q^{s+1-j_s}-1}{q-1}, \text{ where } j_s = \max\{j \in [1,r] \,|\, \lambda^{\rho}_{\dot{n}_s} \ge \mu^{\rho}_{\dot{n}'_j}\} \text{ for } s \in [1,r]. \tag{3.48$'$}$$

3.6. **The q-length function of $S^{\lambda,\mu}$.** Elements $\sigma \in S^{\lambda,\mu}$ can be described in the following way (cf. the proof of Proposition 3.18 and Example 3.20):

Description 3.19. For $s = r, ..., 1$, each $\sigma(s)$ can be any one of the numbers $r, r-1, ..., i_s$ which has not yet been occupied by $\sigma(t)$ for some $t > s$ (cf. (3.50)), with an additional condition that if $\widehat{c}_{a,b}(\lambda) = 1$ for some $a < b$ such that b has not yet been chosen, then $\sigma(s) \ne a$.

We can associate each $\sigma \in S^{\lambda,\mu}$ with a graph defined as follows: Put r weighted points at the bottom such that the s-th point (which will be referred to as *point s^-*) has weight $\mu^{\rho}_{\dot{n}'_s}$. Similarly we put r weighted points on the top such that the s-th point (which will be referred to as *point s^+*) has weight $\lambda^{\rho}_{\dot{n}_s}$. Two points s^+ and t^+ on the top are connected by a line if and only if $\widehat{c}_{s,t}(\lambda) = 1$, in this case we say that the two points are *linked.* Note that if s^+ is linked to t^+, then s^+ is linked to p^+ for all p with

$s<p<t$ (cf. (3.31)). Each $\sigma\in Sym_r$ can be represented by a graph which is obtained by drawing a line (denoted by $L(s)$) between the bottom point s^- and the top point $\sigma(s)^+$ for each $s\in[1,r]$. The length $\ell(\sigma)$ is simply the number of crossings  in the graph. Then $\sigma\in S^{\lambda,\mu}$ if and only if the weight of $\sigma(s)^+$ is not less than the weight of s^- for each s, and in the case $L(t)$ crosses $L(s)$, the two points $\sigma(s)^+$ and $\sigma(t)^+$ on the top cannot be linked, i.e., a graph with the part  is not allowed. See the example below.

Example 3.20. Let λ be given in (3.8) and

$$\mu=\big(\,7,\ 4,\ \overset{4}{4},\ \overset{3}{4},2,1,\overset{2}{1},\overset{1}{1},0\,\big|\,1,\ \overset{1}{1},\overset{2}{1},2,4,\ \overset{3}{4},\ \overset{4}{4},\ 7\big),\quad\text{and so}\tag{3.51}$$

$$\mu^\rho=\big(16,12,\underline{11},\underline{10},7,5,\underline{4},\underline{3},1\ \big|\ 2,\underline{3},\underline{4},6,9,\underline{10},\underline{11},15\big)\qquad\text{(cf. (3.9))}.\tag{3.52}$$

Thus $\mu\preccurlyeq\lambda$. The elements of $\widetilde{S}^{\lambda,\mu}$ correspond to the following graphs:



We see that $i_4=i_3=3$, $i_2=i_1=1$ (where i_s's are defined in (3.49)), thus $\widetilde{S}^{\lambda,\mu}(q)=(1+q^{4-3})1(1+q^{2-1})1=(1+q)^2$, which agrees with the above graphs.

The elements of $S^{\lambda,\mu}$ are represented by the first two of the above graphs. Thus

$$S^{\lambda,\mu}(q)=1+q.$$

Now let us compute the q-length function $S^{\lambda,\mu}(q)$. First we introduce a family of q-functions $Z_q(x;b)$ defined on the set of pairs (x,b) of lexical r-tuples $x=(x_1,...,x_r)$, $b=(b_1,...,b_r)\in\mathbb{Z}^r$ satisfying $1\le b_s\le s$ for $s\in[1,r]$, i.e.,

$$x_1\le x_2\le...\le x_r,\quad b_1\le b_2\le\cdots\le b_r\ \text{ and }\ 1\le b_s\le s,\ \forall\, s\in[1,r].\tag{3.53}$$

Definition 3.21. Define the q-function $Z_q(x;b)$ as follows: Set $Z_q(x;b)=0$ if (3.53) is not satisfied, and define $Z_q(x;b)$ inductively on r by:

$$Z_q(x_1;1)=1,\qquad Z_q(x_1,x_2;1,b_2)=1+\theta(x_2-x_1-1)\theta(1-b_2)q.\tag{3.54}$$

$$\begin{aligned}Z_q(x;b)&=Z_q(x_1,...,x_{r-1};b^{(r-1)})\\&\quad+\sum_{i=b_r}^{r-1}\theta(x_{i+1}-x_i-1)Z_q(x_1,...,x_{i-1},x_{i+1}-1,...,x_r-1;b^{(r-1)})q^{r-i},\end{aligned}\tag{3.55}$$

where $b^{(r-1)}=(b_1,...,b_{r-1})$, and $\theta(x)$ is the *step function* defined by

$$\theta(x)=\begin{cases}1 & \text{if }\ x\ge0,\\ 0 & \text{otherwise.}\end{cases}\tag{3.56}$$

Note that there are only finite many functions $Z_q(x;b)$ for each fixed r. To see this, for any lexical $x\in\mathbb{Z}^r$ and for $1\le s\le t\le r$, we define $c_{s,t}(x)$ and $\widehat{c}_{s,t}(x)$ analogous to (3.26) and (3.27) by

$$c_{s,t}(x)=\begin{cases}1 & \text{if }\ t=s\text{ or }x_t-x_s<t-s,\\ 0 & \text{otherwise,}\end{cases}\tag{3.57}$$

$$\widehat{c}_{s,t}(x)=\begin{cases}1 & \text{if }\ c_{s,s}(x)=c_{s,s+1}(x)=...=c_{s,t}(x)=1,\\ 0 & \text{otherwise,}\end{cases}\tag{3.58}$$

and define $\widehat{x} = (\widehat{x}_1, ..., \widehat{x}_r)$ with

$$\widehat{x}_s = \#\{p \in [1, s-1] \mid \widehat{c}_{p,s}(x) = 0\} \quad \text{for } \ s \in [1, r]. \tag{3.59}$$

Then one can prove that $\widehat{x}$ is lexical and $0 \le \widehat{x}_s < s$ for $s \in [1, r]$, and

$$Z_q(x; b) = Z_q(\widehat{x}; b). \tag{3.60}$$

Thus there are only finite many functions $Z_q(x, b)$ for each fixed r.

Proposition 3.22. *Let $\lambda, \mu \in P_r^{\mathrm{Lex}}$ with $\mu \preccurlyeq \lambda$. Denote*

$$b^{\lambda,\mu} = (i_1, ..., i_r) \in \mathbb{Z}^r, \ \textit{where } i_s = \min\{i \in [1, r] \,|\, \lambda^\rho_{\dot{n}_s} \le \mu^\rho_{\dot{n}'_i}\} \ \ \textit{for } s \in [1, r], \tag{3.61}$$

namely i_s is defined in (3.49). Let $h(\lambda)$ be the height vector of λ defined in (3.7). Then the q-length function of $S^{\lambda,\mu}$ is given by

$$S^{\lambda,\mu}(q) = Z_q(h(\lambda); b^{\lambda,\mu}). \tag{3.62}$$

Proof. The proposition follows from Description 3.19 and Definition 3.6. ☐

3.7. **Generalized Kazhdan-Lusztig polynomials.** Let $K_{\lambda,\mu}(q)$ be the Kazhdan-Lusztig polynomial defined in [12] (which was denoted as $K_{\mu,\lambda}(q)$ in [2], but we prefer its original notation in [12]), and $\ell_{\lambda,\mu}(q)$ the polynomial defined in [2] (which was denoted as $\ell_{\mu,\lambda}(q)$ in [2]). Then [2, Corollary 3.39] states that

$$K_{\lambda,\mu}(q) = \ell_{\lambda,\mu}(-q^{-1}) = \sum_{\theta \in \mathbb{N}^{\#\lambda}:\, R'_\theta(\mu) = \lambda} q^{|\theta|}, \quad \text{where } |\theta| = \sum_{s=1}^r \theta_s, \tag{3.63}$$

for dominant weights λ, μ. Let $\ell(\lambda, \mu)$ be the length defined in (3.17). First let us see an example of computing $\ell(\lambda, \mu)$.

Example 3.23. Suppose λ is given in (3.8) and μ given in (3.51). By (3.16),

$$\ell_1(\lambda, \mu) = 2 - 1 + 2 - 2 = 1, \qquad \ell_2(\lambda, \mu) = 3 - 1 + 3 - 4 = 1,$$
$$\ell_3(\lambda, \mu) = 5 - 4 + 6 - 6 = 1, \qquad \ell_4(\lambda, \mu) = 6 - 4 + 7 - 7 = 2.$$

Thus $\ell(\lambda, \mu) = 7$.

One of the main results of this paper is the following.

Theorem 3.24. *Suppose λ, μ are dominant weights. Then $K_{\lambda,\mu}(q) \ne 0$ if and only if $\mu \preccurlyeq \lambda$ and in this case*

$$K_{\lambda,\mu}(q) = q^{\ell(\lambda,\mu)} \sum_{\sigma \in S^{\lambda,\mu}} q^{-2\ell(\sigma)} = q^{\ell(\lambda,\mu)} S^{\lambda,\mu}(q^{-2}), \tag{3.64}$$

where $S^{\lambda,\mu}$ is defined by (3.41), and $S^{\lambda,\mu}(q)$ is the q-length function of $S^{\lambda,\mu}$ defined in (3.45) and can be determined by (3.62).

A weight λ is said to be

- *totally disconnected* if $c_{s,t}(\lambda) = 0$ for all pairs (s, t) with $s < t$;
- *totally connected* if $c_{s,t}(\lambda) = 1$ for all pairs (s, t) with $s \le t$.

From Theorem 3.24, one immediately obtains

Corollary 3.25. *Let $\lambda, \mu \in P_r^{\mathrm{Lex}}$ with $\mu \preccurlyeq \lambda$, and let $S^\lambda(q)$, $\widetilde{S}^{\lambda,\mu}(q)$ be as in Proposition 3.18. We have*

1. *If λ is totally connected, then $K_{\lambda,\mu}(q) = q^{\ell(\lambda,\mu)}$.*
2. *If λ is totally disconnected, then $K_{\lambda,\mu}(q) = q^{\ell(\lambda,\mu)} \widetilde{S}^{\lambda,\mu}(q^{-2})$.*
3. *If $\mu \prec\!\prec \lambda$ (recall (3.37)), then $K_{\lambda,\mu}(q) = q^{\ell(\lambda,\mu)} S^\lambda(q^{-2})$.*

3.8. **Proof of Theorem 3.24.** Using notations as above, we set (cf. (3.63))

$$\Theta^\lambda_\mu=\{\theta\in\mathbb{N}^r\,|\,R'_\theta(\mu)=\lambda\},\qquad\text{where }\ r=\#\lambda.$$

In the three lemmas below, we shall establish a bijection between Θ^λ_μ and $S^{\lambda,\mu}$. Theorem 3.24 is then a simple consequence of this bijection.

First let us define a map

$$\Theta^\lambda_\mu\longrightarrow\widetilde{S}^{\lambda,\mu}:\ \theta\mapsto\sigma_\theta \tag{3.65}$$

in the following way. Suppose $\theta\in\Theta^\lambda_\mu$, i.e. (cf. (3.15) for notations m'_s,n'_s),

$$\lambda=R'_\theta(\mu)=(R^{\theta_1}_{m'_1,\dot n'_1}\circ\cdots\circ R^{\theta_r}_{m'_r,\dot n'_r}(\mu))^+. \tag{3.66}$$

We denote

$$\mu^{(r)}=\mu,\ \ \mu^{(s-1)}=R^{\theta_s}_{m'_s,\dot n'_s}(\mu^{(s)})\qquad\text{for }\ s=r,r-1,...,1. \tag{3.67}$$

Then each $\mu^{(s-1)}$ is obtained from $\mu^{(s)}$ by adding some number, say, K_s, to its m'_s-th, $\dot n'_s$-th entries. In fact (we use the notation $k^{\mu^{(s,i)}}_s$ to denote the integer k_s defined in (3.29) with λ replaced by $\mu^{(s,i)}$)

$$K_s=\sum_{i=1}^{\theta_s}k^{\mu^{(s,i)}}_s,\quad\text{where }\mu^{(s,0)}=\mu^{(s)},\ \mu^{(s,i)}=R_{m'_s,\dot n'_s}(\mu^{(s,i-1)}). \tag{3.68}$$

Thus

$$\mu^{(0)}=\mu+\sum_{s=1}^{r}K_s\gamma'_s. \tag{3.69}$$

Since $\lambda=(\mu^{(0)})^+$ and both λ and $\mu^{(0)}$ are regular, there exists a $\sigma\in Sym_r$ uniquely determined by θ, such that (cf. (3.38) and (3.39))

$$\text{aty}_{\mu^{(0)}}=\sigma(\text{aty}_\lambda)=\text{aty}_{\sigma\cdot\lambda},\quad\text{i.e., }\ (\mu^{(0)})^\rho_{\dot n_s}=\lambda^\rho_{\dot n_{\sigma(s)}}\ \text{ for }s\in[1,r]\ \text{ (cf. (3.4))}. \tag{3.70}$$

By (3.69), $\text{aty}_\mu\le\text{aty}_{\mu^{(0)}}=\text{aty}_{\sigma\cdot\lambda}$, and by (3.13) and (3.36), $\text{typ}_\mu=\text{typ}_\lambda$. But $\text{typ}_\lambda=\text{typ}_{\sigma\cdot\lambda}$ (typical tuples are invariant under the dot action of Sym_r, cf. (3.5), (3.38) and (3.39)). Thus $\mu\preccurlyeq\sigma\cdot\lambda$ by (3.13), i.e., $\sigma\in\widetilde{S}^{\lambda,\mu}$ by definition (3.42). We denote σ by σ_θ. Thus we obtain the map (3.65).

Lemma 3.26. *The map (3.65) is an injection. More precisely, suppose $\sigma\in Sym_r$ such that $\sigma=\sigma_\theta$ for some $\theta=(\theta_1,...,\theta_r)\in\Theta^\lambda_\mu$, then such θ is unique and is given by*

$$\theta_s=\theta'_s-2\ell(\sigma,s)\qquad\textit{(cf. (3.44)), where} \tag{3.71}$$

$$\theta'_s=\#Q_s,\quad Q_s=[\mu^\rho_{\dot n'_s}+1,\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\text{Set}(\text{typ}_\lambda)\qquad\textit{for }\ s=r,r-1,...,1. \tag{3.72}$$

Thus by (3.22)), θ'_s is in fact $\theta'_s=h_{\sigma(s)}(\lambda)-h_s(\mu)+\sigma(s)-s$.

Proof. The proof of the lemma is divided into the following cases.

Case 1: $s=r$.

We want to prove $\theta_r=\theta'_r$. Note that each time when we apply $R_{m'_r,\dot n'_r}$ to μ, the r-th entry $\mu^\rho_{\dot n'_r}$ of the atypical tuple aty_μ reaches an integer in the set Q_r (cf. (3.32) and (3.33)), and no integer in this set Q_r can be skipped. Thus after applying θ'_r times, this entry reaches the integer $\lambda^\rho_{\dot n_{\sigma(r)}}$ (it is an entry of aty_λ, thus not in the typical entry set $\text{Set}(\text{typ}_\lambda)$), and μ becomes $\mu^{(r-1)}$ (cf. (3.67)). Thus $\theta_r=\theta'_r$.

One can also use the following arguments to prove $\theta_r=\theta'_r$: The integer $\lambda^\rho_{\dot n_{\sigma(r)}}$ is the r-th entry of the atypical tuple $\text{aty}_{\mu^{(r-1)}}$, which by definitions (3.33) and (3.35) and by (3.32), is equal to the θ_r-th smallest integer bigger than $\mu^\rho_{\dot n'_r}$ and not in $\text{Set}(\text{typ}_\lambda)$. But by the definition of θ'_r in (3.72), $\lambda^\rho_{\dot n_{\sigma(s)}}$ is the θ'_r-th smallest integer bigger than $\mu^\rho_{\dot n'_r}$ and not in Set(typ$_\lambda$). Thus $\theta_r = \theta'_r$.

Case 2: $s = r-1$ and $\sigma(r-1) < \sigma(r)$.

We want to prove $\theta_{r-1} = \theta'_{r-1}$. There are two possibilities to consider.

Subcase 2.i: Suppose $\mu^\rho_{\dot n'_{r-1}} = \lambda^\rho_{\dot n_{\sigma(r-1)}}$. Then $\mu^{(r-2)} = \mu^{(r-1)}$ and we obviously have $\theta_{r-1} = 0 = \theta'_{r-1}$. We are done.

Subcase 2.ii: Suppose $\mu^\rho_{\dot n'_{r-1}} < \lambda^\rho_{\dot n_{\sigma(r-1)}}$. Note that $\lambda^\rho_{\dot n_{\sigma(r-1)}} < \lambda^\rho_{\dot n_{\sigma(r)}}$. Also note that

$$\mathrm{Set}((\mu^{(r-1)})^\rho) = (\mathrm{Set}(\mu^\rho)\backslash\{\mu^\rho_{\dot n'_r}\}) \cup \{\lambda^\rho_{\dot n_{\sigma(r)}}\}, \tag{3.73}$$

(recall the ρ-translated notation in (3.3) and note that the only difference between $\mu^{(r-1)}$ and μ is their r-th atypical entries). We observe that $\lambda^\rho_{\dot n_{\sigma(r-1)}}$ is not an entry of the typical tuple typ$_\mu$ = typ$_\lambda$ because of the regularity of λ, it is not the t-th entry of the atypical tuple aty$_\mu$ either for any $t \le r-1$ because of the assumption that $\mu^\rho_{\dot n'_{r-1}} < \lambda^\rho_{\dot n_{\sigma(r-1)}}$. Thus $\lambda^\rho_{\dot n_{\sigma(r-1)}}$ is not in (3.73). But it is in the set

$$[(\mu^{(r-1)})^\rho_{\dot n'_{r-1}}, (\mu^{(r-1)})^\rho_{\dot n'_r}] = [\mu^\rho_{\dot n'_{r-1}}, \lambda^\rho_{\dot n_{\sigma(r)}}]. \tag{3.74}$$

Hence there is at least an integer in (3.74) but not in (3.73). By Definition 3.6, the $(r-1)$-th and r-th atypical roots of $\mu^{(r-1)}$ are not c-related.

Similarly, the $(r-1)$-th and r-th atypical roots of $\mu^{(r-1,i)}$ (cf. (3.68)) are not c-related for all i with $1 \le i < \theta'_{r-1}$. Thus by the arguments in Case 1, we need to apply the raising operator $R_{m'_{r-1},\dot n'_{r-1}}$ to $\mu^{(r-1)}$ exactly θ'_{r-1} times in order to obtain $\mu^{(r-2)}$. So $\theta'_{r-1} = \theta_{r-1}$.

Case 3: $s = r-1$ and $\sigma(r-1) > \sigma(r)$.

We want to prove $\theta_{r-1} = \theta'_{r-1} - 2$. Note that

$$(\mu^{(r-1)})^\rho_{\dot n'_{r-1}} = \mu^\rho_{\dot n'_{r-1}} < \mu^\rho_{\dot n'_r} \le (\mu^{(0)})^\rho_{\dot n'_r} = \lambda^\rho_{\dot n_{\sigma(r)}} < \lambda^\rho_{\dot n_{\sigma(r-1)}} \quad \text{(cf. (3.70))}, \tag{3.75}$$

(recall that λ, μ are dominant). Thus we need to apply $R_{m'_{r-1},\dot n'_{r-1}}$ to $\mu^{(r-1)}$ at least once.

Suppose after applying i times of $R_{m'_{r-1},\dot n'_{r-1}}$ to $\mu^{(r-1)}$ (i can be zero), the $(r-1)$-th entry $(\mu^{(r-1)})^\rho_{\dot n'_{r-1}}$ of the atypical tuple aty$_{\mu^{(r-1)}}$ reaches an integer, say p, such that

$$p < \lambda^\rho_{\dot n_{\sigma(r)}} \quad \text{but} \quad [p, \lambda^\rho_{\dot n_{\sigma(r)}}] \subset \mathrm{Set}(\lambda^\rho). \tag{3.76}$$

By (3.75), such i must exist since the $(r-1)$-th entry will finally reach the integer $\lambda^\rho_{\dot n_{\sigma(r-1)}}$.

Note that p and $\lambda^\rho_{\dot n_{\sigma(r)}}$ are respectively the $(r-1)$-th and r-th entries of aty$_{\mu^{(r-1,i)}}$ (cf. (3.68)). Thus by (3.76) and Definition 3.6, the $(r-1)$-th and r-th atypical roots of $\mu^{(r-1,i)}$ are c-related. Then by (3.32) and (3.33), when we apply $R_{m'_{r-1},\dot n'_{r-1}}$ to $\mu^{(r-1,i)}$, the $(r-1)$-th entry of aty$_{\mu^{(r-1,i)}}$ reaches an integer in the set

$$[\lambda^\rho_{\dot n_{\sigma(r)}} + 1, \lambda^\rho_{\dot n_{\sigma(r-1)}}]\backslash\mathrm{Set}(\mathrm{typ}_{\mu^{(r-1,i)}}) = [\lambda^\rho_{\dot n_{\sigma(r)}} + 1, \lambda^\rho_{\dot n_{\sigma(r-1)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda), \tag{3.77}$$

(recall that Set(typ$_{\mu^{(r-1,i)}}$) = Set(typ$_\lambda$)), such that an integer in this set is skipped. So

$$\#\left([\lambda^\rho_{\dot n_{\sigma(r)}}, \lambda^\rho_{\dot n_{\sigma(r-1)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda)\right) \ge 2. \tag{3.78}$$

In particular there is at least an element in the set (3.77), which means

$$\widehat{c}_{\sigma(r),\sigma(r-1)}(\lambda)=0 \quad \text{if} \ \ \sigma(r-1)-\sigma(r)=1 \quad \text{(cf. (3.26))}. \tag{3.79}$$

Note that $\lambda^\rho_{\dot n_{\sigma(r)}}$ is in $Q_{r-1}=[\mu^\rho_{\dot n'_{r-1}}+1,\lambda^\rho_{\dot n_{\sigma(r-1)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda)$ by (3.75), but not in the set (3.77). Thus we have in fact skipped two integers in the set Q_{r-1}. Therefore

$$\theta_{r-1}=\theta'_{r-1}-2 \qquad \text{if} \quad \sigma(r-1)>\sigma(r).$$

Case 4: The general case.

In general, when we apply $R_{m'_s,\dot n'_s}$ to $\mu^{(s)}$ in order to obtain $\mu^{(s-1)}$, for each $t>s$ with $\sigma(t)<\sigma(s)$ the above arguments show that we have to skip two integers in the set Q_s. Therefore we have (3.71) and the lemma. □

Lemma 3.27. *The image of the map* (*3.65*) *is contained in* $S^{\lambda,\mu}$, *i.e., if* $\sigma=\sigma_\theta$ *for some* $\theta\in\Theta^\lambda_\mu$, *then*

$$\widehat{c}_{\sigma(t),\sigma(s)}(\lambda)=0 \qquad \textit{if} \quad \sigma(t)<\sigma(s) \quad \textit{for any} \quad t>s. \tag{3.80}$$

Proof. Let p be the number such that $\sigma(p)$ is minimal among those $\sigma(u)$ with $u<t$ and $\sigma(u)>\sigma(t)$. Then $p=s$ or $\sigma(p)<\sigma(s)$. The definition (3.27) means that the relation $\widehat{c}_{\sigma(t),\sigma(p)}(\lambda)=0$ implies the relation (3.80). Thus it suffices to prove $\widehat{c}_{\sigma(t),\sigma(p)}(\lambda)=0$.

For any p' with $\sigma(t)\le\sigma(p')<\sigma(p)$, by the choice of p, we have $p'\ge t$ and so $p'>p$. Thus the arguments in the proof of Lemma 3.26 show that when we apply $R_{m'_p,\dot n'_p}$ to $\mu^{(p)}$ in order to obtain $\mu^{(p-1)}$, we need to pass over the integer $\lambda^\rho_{\dot n_{\sigma(p')}}$ and skip another integer for all such p'. Hence there are at least $2(\sigma(p)-\sigma(t))$ integers in the set $[\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(p)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda)$ (cf. (3.78)). This means that there are at least $\sigma(p)-\sigma(t)$ integers in the set $[\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(p)}}]\backslash\mathrm{Set}(\lambda^\rho)$ (since there are exactly $\sigma(p)-\sigma(t)$ integers in $[\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(p)}}]$ which are entries of atypical tuple aty_λ), which implies that $c_{\sigma(t),\sigma(p)}(\lambda)=0$ by (3.26) and so $\widehat{c}_{\sigma(t),\sigma(p)}(\lambda)=0$ (cf. (3.79)). □

Lemma 3.28. *The map* (*3.65*) *is a bijection between* Θ^λ_μ *and* $S^{\lambda,\mu}$.

Proof. For any $\sigma\in S^{\lambda,\mu}$, we define θ_s as in (3.71). We want to prove

$$\theta_s\ge 0 \qquad \text{for} \quad s=1,...,r. \tag{3.81}$$

Suppose $t>s$ such that $\sigma(t)<\sigma(s)$. So

$$\widehat{c}_{\sigma(t),\sigma(s)}(\lambda)=0. \tag{3.82}$$

Denote

$$X_{s,t}=\{p>s\,|\,\sigma(t)\le\sigma(p)<\sigma(s)\} \quad \text{and} \quad x_{s,t}=\#X_{s,t}. \tag{3.83}$$

First we prove by induction on $\sigma(s)-\sigma(t)$ that

$$\#\left([\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\mathrm{Set}(\lambda^\rho)\right)\ge x_{s,t}. \tag{3.84}$$

If $\sigma(s)-\sigma(t)=1$, then (3.84) follows from (3.82) and Definition 3.6 (note that obviously $\sigma(s)-\sigma(t)\ge x_{s,t}$).

In general set $p'=x_{s,t}$ and we write the set $\{\sigma(p)\,|\,p\in X_{s,t}\}$ in ascending order:

$$\{\sigma(p)\,|\,p\in X_{s,t}\}=\{\sigma(t_1)<\sigma(t_2)<\cdots<\sigma(t_{p'})\}, \tag{3.85}$$

where $t_1=t$ and $\sigma(t_{p'})<\sigma(s)$. Since $\widehat{c}_{\sigma(t),\sigma(s)}(\lambda)=0$, by Definition 3.6, there exists some number, denoted by $\sigma(i)$, lies in between $\sigma(t)$ and $\sigma(s)$, i.e.,

$$\sigma(t)<\sigma(i)\le\sigma(s), \tag{3.86}$$

such that $c_{\sigma(t),\sigma(i)}(\lambda)=0$, that is,

$$\#\left([\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(i)}}]\backslash\mathrm{Set}(\lambda^\rho)\right)\geq\sigma(i)-\sigma(t)\quad\text{(cf. (3.19))}.\tag{3.87}$$

Now we prove (3.84) by induction on $\sigma(s)-\sigma(t)$ in two cases.

Case 1: Suppose $i\in X_{s,t}$*, say* $i=t_{p''}$ *for some* $1<p''\leq p'$. Then

$$\#\left([\lambda^\rho_{\dot n_{\sigma(t_{p''})}},\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\mathrm{Set}(\lambda^\rho)\right)\geq x_{s,t_{p''}}\geq p'-p''+1,\tag{3.88}$$

where the first inequality follows from the inductive assumption that (3.84) holds for $t_{p''}$ since $\sigma(s)-\sigma(t_{p''})<\sigma(s)-\sigma(t)$, and the second from the fact that $t_{p''},t_{p''+1},...,t_{p'}\in X_{s,t_{p''}}$. Then (3.84) follows from (3.88) and (3.87) (with i replaced by $t_{p''}$) by noting that $\sigma(t_{p''})-\sigma(t)\geq p''-1$ (cf. (3.85)) and that $p'=x_{s,t}$.

Case 2: Suppose $i\notin X_{s,t}$. This means that $i<s$. Let p'' be the minimal integer with $1\leq p''\leq p'$ such that $\sigma(t_{p''})>\sigma(i)$. Then $\sigma(s)-\sigma(t_{p''})<\sigma(s)-\sigma(t)$ and so (3.88) holds again in this case by the inductive assumption. Furthermore since $\sigma(i)-\sigma(t)<\sigma(s)-\sigma(t)$ (cf. (3.86)), the inductive assumption also gives

$$\#\left([\lambda^\rho_{\dot n_{\sigma(t)}},\lambda^\rho_{\dot n_{\sigma(i)}}]\backslash\mathrm{Set}(\lambda^\rho)\right)\geq x_{i,t}\geq p''-1,\tag{3.89}$$

where the last inequality follows from the fact that

$$t_1,...,t_{p''-1}\in\{p>i\,|\,\sigma(t)\leq\sigma(p)\leq\sigma(i)\}=X_{i,t},$$

(recall that $i<s$). Now (3.84) follows from (3.89) and (3.88). This completes the proof of (3.84).

Now set $p=\ell(\sigma,s)$ and write

$$\{t>s\,|\,\sigma(t)<\sigma(s)\}=\{s_1,s_2,...,s_p\,|\,\sigma(s_1)<\sigma(s_2)<\cdots<\sigma(s_p)<\sigma(s)\}.\tag{3.90}$$

(Then the left-hand side of (3.90) is in fact the set X_{s,s_1}, cf. (3.85).) Thus (3.84) means that the set

$$[\lambda^\rho_{\dot n_{\sigma(s_1)}},\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\mathrm{Set}(\lambda^\rho)\tag{3.91}$$

has cardinality $\geq x_{s,s_1}=p$, and so the set

$$[\mu^\rho_{\dot n'_s}+1,\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda)\tag{3.92}$$

has cardinality $\geq 2p$, because $\mu^\rho_{\dot n'_s}<\mu^\rho_{\dot n'_{s_i}}\leq\lambda^\rho_{\dot n_{\sigma(s_i)}}<\lambda^\rho_{\dot n_{\sigma(s)}}$, i.e., we have p more elements $\lambda^\rho_{\dot n_{\sigma(s_i)}}$ with $i\in[1,p]$ which are not in the first set (3.91) but in the second set (3.92).

Then (3.72), (3.44) and the fact that (3.92) has cardinality $\geq 2p$ show that $\theta'_s\geq 2p$ and so $\theta_s\geq 0$. (In fact if $p>0$ then $\theta_s\geq 1$ since in this case there exists at least one more integer $\lambda^\rho_{\dot n_{\sigma(s)}}$ which is in $[\mu^\rho_{\dot n'_s}+1,\lambda^\rho_{\dot n_{\sigma(s)}}]\backslash\mathrm{Set}(\mathrm{typ}_\lambda)$.) This proves (3.81).

Now we define $\mu^{(0)}$ as in (3.67). Then the arguments in the proof of Lemma 3.26 show that (3.70) holds, i.e., $(\mu^{(0)})^+=\lambda$. Thus $\theta\in\Theta^\lambda_\mu$ and $\sigma=\sigma_\theta$. Therefore $\theta\mapsto\sigma_\theta$ is a bijection between Θ^λ_μ and $S^{\lambda,\mu}$. □

Proof of Theorem 3.24. Finally we return to the proof of Theorem 3.24. By (3.17), (3.23), (3.43) and (3.72), we have

$$|\theta|=\sum_{s=1}^r\theta'_s-\sum_{s=1}^r\ell(\sigma,s)=\ell(\lambda,\mu)-\ell(\sigma).$$

Now (3.64) follows from (3.63). □

3.9. **A correspondence between r-fold atypical modules over $\mathfrak{gl}_{m|n}$ and $\mathfrak{gl}_{r|r}$.** A subset $\mathcal{B} \subset \mathbb{Z}_+^{m|n}$ of dominant integral weights is called a *block of* $\mathbb{Z}_+^{m|n}$ *for* $\mathfrak{gl}_{m|n}$ if it is a maximal subset such that for any two weights $\lambda, \mu \in \mathcal{B}$, there exist weights $\mu^1, \mu^2, ..., \mu^k \in \mathcal{B}$ with $\lambda = \mu^1$, $\mu^k = \mu$ such that the extension group $\mathrm{Ext}^1(V(\mu^i), V(\mu^{i+1})) \neq 0$ for $i = 1, ..., k-1$. Then $\mathbb{Z}_+^{m|n}$ is divided into a disjoint union of blocks. Lemma 1.12 in [11] says that for any $\lambda, \mu \in \mathcal{B}$, one has $\#\lambda = \#\mu$, which is called the *degree of atypical type of* $\mathcal{B}$, and denoted by $\#\mathcal{B}$. Let $\mathcal{E}^{m|n}$ be the category of finite dimensional $\mathfrak{gl}_{m|n}$-modules. A dominant weight μ is called a *primitive weight of a module* V if it is the highest weight of a composition factor of V. Denote by $\mathrm{Prim}(V)$ the set of primitive weights of V. For $\lambda, \mu \in \mathbb{Z}_+^{m|n}$, we denote by $a_{\lambda,\mu} = [\overline{V}(\lambda) : V(\mu)]$ the multiplicity of the composition factor $V(\mu)$ in the Kac-module $\overline{V}(\lambda)$. It was proved in [2] that

$$a_{\lambda,\mu} \le 1, \quad \text{and the matrix } (a_{\lambda,\mu}) \text{ is the inverse of the matrix } (K_{\lambda,\mu}(-1)), \tag{3.93}$$

where the matrices were defined with respect to some total order of weights compatible with the partial order "$\preccurlyeq$".

An application of Theorem 3.24 is the following.

Theorem 3.29. *Let* $\mathcal{B}$ *be a block of* $\mathbb{Z}_+^{m|n}$ *for* $\mathfrak{gl}_{m|n}$ *with* $\#\mathcal{B} = r$. *Let* $\mathcal{B}'$ *be the unique block of* $\mathbb{Z}_+^{r|r}$ *for* $\mathfrak{gl}_{r|r}$ *with* $\#\mathcal{B}' = r$.

(1) *There exists a bijection*

$$\phi : \mathcal{B} \to \mathcal{B}', \qquad \phi(\lambda) = (h'(\lambda) \,|\, h(\lambda)), \tag{3.94}$$

where $h(\lambda)$ *is the height vector of* λ *defined by (3.7) and* $h'(\lambda) = (h_r(\lambda), ..., h_1(\lambda))$.

(2) *The Kazhdan-Lusztig polynomials* $K_{\lambda,\mu}(q)$ *of* $\mathfrak{gl}_{m|n}$ *and* $K_{\phi(\lambda),\phi(\mu)}(q)$ *of* $\mathfrak{gl}_{r|r}$ *coincide, that is,*

$$K_{\lambda,\mu}(q) = K_{\phi(\lambda),\phi(\mu)}(q) \tag{3.95}$$

for $\lambda, \mu \in \mathcal{B}$.

(3) *Under the mapping (3.94), the set* $\mathrm{Prim}(\overline{V}(\lambda))$ *of primitive weights of the* $\mathfrak{gl}_{m|n}$*-Kac-module* $\overline{V}(\lambda)$ *for* $\lambda \in \mathcal{B}$ *is in one to one correspondence to the set* $\mathrm{Prim}(\overline{V}(\phi(\lambda)))$ *of primitive weights of the* $\mathfrak{gl}_{r|r}$*-Kac-module* $\overline{V}(\phi(\lambda))$, *namely*

$$\mathrm{Prim}(\overline{V}(\phi(\lambda))) = \{\phi(\mu) \,|\, \mu \in \mathrm{Prim}(\overline{V}(\lambda))\}. \tag{3.96}$$

Proof. (1) From Theorem 3.24 (cf. (3.36)) we can deduce that any two dominant weights are in the same block if and only if their typical tuples are equal (thus blocks of $\mathbb{Z}_+^{m|n}$ are in one to one correspondence with the typical tuples). Therefore for any block $\mathcal{B}$ of $\mathbb{Z}_+^{m|n}$ with $\#\mathcal{B} = r$, we can denote the typical tuple of any weight in $\mathcal{B}$ by

$$\mathrm{tt}_{\mathcal{B}} = (\mathrm{tt}_1, ..., \mathrm{tt}_{m-r} \,|\, \mathrm{tt}_{\dot{r}+1}, ..., \mathrm{tt}_{\dot{n}}), \tag{3.97}$$

where $\mathrm{tt}_1 > ... > \mathrm{tt}_{m-r}$, $\mathrm{tt}_{\dot{r}+1} < ... < \mathrm{tt}_{\dot{n}}$ are all distinct.

First we prove that the map ϕ is injective, which is equivalent to proving that for any two dominant weights $\lambda, \mu \in \mathcal{B}$ with $h(\lambda) = h(\mu)$, one has the equality of the atypical tuples $\mathrm{aty}_\lambda = \mathrm{aty}_\mu$. Thus assume that $\lambda^\rho_{\dot{n}_s} \neq \mu^\rho_{\dot{n}'_s}$ for some s. Say, $\lambda^\rho_{\dot{n}_s} > \mu^\rho_{\dot{n}'_s}$. Then the right-hand side of (3.22) with $t = s$ is zero, but the left-hand side is not since $\lambda^\rho_{\dot{n}_s} \in [\mu^\rho_{\dot{n}'_s} + 1, \lambda^\rho_{\dot{n}_s}] \backslash \mathrm{Set}(\mathrm{tt}_{\mathcal{B}})$.

Next we prove that the map ϕ is surjective, which is equivalent to proving that for any lexical r-tuple $a = (a_1, ..., a_r) \in \mathbb{Z}^r$, there exists $\lambda \in \mathcal{B}$ such that $h(\lambda) = a$. Fix a lexical r-tuple $b = (b_1, ..., b_r) \in \mathbb{Z}^r$ such that $b_r < \min\{\mathrm{tt}_{m-r}, \mathrm{tt}_{\dot{r}+1}, a_1\} - r$. Let

$$\mu = -\rho + (\mathrm{tt}_1, ..., \mathrm{tt}_{m-r}, b_r + r, ..., b_1 + 1 \,|\, b_1 + 1, ..., b_r + r, \mathrm{tt}_{\dot{r}+1}, ..., \mathrm{tt}_{\dot{n}}) \in \mathbb{Z}^{m|n}. \quad (3.98)$$

Then μ is dominant such that $\mathrm{typ}_\mu = \mathrm{tt}_{\mathcal{B}}$ (thus $\mu \in \mathcal{B}$) and $h(\mu) = b$. Take $t = s$ and $h_s(\lambda) = a_s$ in (3.22), then (3.22) uniquely determines a number $\lambda^\rho_{\dot{n}_s} \notin \mathrm{Set}(\mathrm{tt}_{\mathcal{B}})$. This uniquely determines a weight $\lambda \in \mathcal{B}$. To see $h(\lambda) = a$, use (3.22) again. Thus ϕ is a bijection.

In fact the atypical tuple aty_λ determines the height vector $h(\lambda)$ through the following formula:

$$h_s(\lambda) + s = \lambda^\rho_{\dot{n}_s} - \#\{\text{typical entries which are smaller than } \lambda^\rho_{\dot{n}_s}\}. \quad (3.99)$$

Conversely the height vector $h(\lambda)$ determines the atypical tuple aty_λ as follows: First set $\mathrm{aty}_\lambda = h(\lambda)$, and denote aty_λ as $\mathrm{aty}_\lambda = (a_1, ..., a_r)$. Label the typical entries of typ_λ in ascending order: $x_1 < x_2 < ... < x_{m+n-r}$. For each $i = 1, 2, ..., m+n-r$, if $a_t \geq x_i$ then replace a_t by $a_t + 1$ for all $t = 1, 2, ..., r$.

(2) Note that in $\mathfrak{gl}_{r|r}$, we have $\mathrm{aty}_{\phi(\lambda)} = (\phi(\lambda))^{\rho_{r|r}}$ (cf. notation (3.3)), where $\rho_{r|r} = (r, ..., 1 \,|\, 1, ..., r)$, and the typical tuple $\mathrm{typ}_{\phi(\lambda)}$ is empty. Also we have the equality of the height vectors: $h(\lambda) = h(\phi(\lambda))$. Thus by Definition 3.6, we have $c_{s,t}(\lambda) = c_{s,t}(\phi(\lambda))$ for all $s, t \in [1, r]$. Let μ be another dominant weight with $\mu \preccurlyeq \lambda$. By (3.22) and (3.61), we have $b^{\lambda,\mu} = b^{\phi(\lambda),\phi(\mu)}$. This gives (3.95) by (3.60) and (3.64).

(3) Using (3.93) and (3.95), we have

$$\begin{aligned} \mathrm{Prim}(\overline{V}(\phi(\lambda))) &= \{\mu' \,|\, a_{\phi(\lambda),\mu'} = 1\} \\ &= \{\phi(\mu) \,|\, a_{\phi(\lambda),\phi(\mu)} = 1\} = \{\phi(\mu) \,|\, a_{\lambda,\mu} = 1\} = \phi(\mathrm{Prim}(\overline{V}(\lambda))). \qquad \square \end{aligned}$$

4. Character formulae

As mentioned in the introduction, this section contains three main results: the proof of the conjecture due to van der Jeugt et al, the construction of a Kac-Weyl type character formula, and the derivation of a dimension formula.

We shall continue to use notations in the previous sections. Moreover, we define

$$m(\Lambda)_\lambda = \# S^{\Lambda,\lambda} \quad (4.1)$$

to be the cardinality of the set $S^{\Lambda,\lambda}$ (cf. (3.40)). Then $m(\Lambda)_\lambda = S^{\Lambda,\lambda}(1)$ (cf. (3.45)).

For any weight $\lambda \in P$, we define what is called the *Kac-character of* λ:

$$\chi^{\mathrm{Kac}}(\lambda) = \frac{L_1}{L_0} \sum_{w \in W} \epsilon(w) e^{w(\lambda+\rho)}. \quad (4.2)$$

Namely, $\chi^{\mathrm{Kac}}(\lambda)$ is defined by the right-hand side of (2.13). Thus it is the character of the Kac-module $\overline{V}(\lambda)$ when λ is dominant.

4.1. Proof of the conjecture of van der Jeugt et al.

As an immediate consequence of Theorem 3.24, we have

Theorem 4.1. *The formal character* $\mathrm{ch}\, V(\Lambda)$ *of the finite dimensional irreducible* $\mathfrak{g}$-*module* $V(\Lambda)$ *is given by*

$$\mathrm{ch}\, V(\Lambda) = \sum_{\lambda \in P_+ :\, \lambda \preccurlyeq \Lambda} (-1)^{\ell(\Lambda,\lambda)} m(\Lambda)_\lambda \chi^{\mathrm{Kac}}(\lambda), \quad (4.3)$$

where P_+ is the set of dominant integral weights, the length $\ell(\Lambda,\lambda)$ is defined in (3.17), and the partial order "$\preccurlyeq$" is defined in (3.13).

Proof. This follows from (3.64), (4.1) and [12, Lemma 3.4], which states

$$\mathrm{ch}\,V(\Lambda)=\sum_{\lambda\in P_+}K_{\Lambda,\lambda}(-1)\chi^{\mathrm{Kac}}(\lambda). \qquad \square$$

One can re-write (4.3) to obtain the conjecture of van der Juegt et al. To state it, we need to introduce the following notations.

Let λ as in (3.1) be an r-fold atypical weight (not necessarily dominant) with atypical roots ordered as in (3.2): $\gamma_1<...<\gamma_r$. We define the *normal cone with vertex* λ:

$$\mathcal{C}_\lambda^{\mathrm{Norm}}=\{\lambda-\sum_{s=1}^{r}i_s\gamma_s\,|\,i_s\geq 0\}. \tag{4.4}$$

We also define $\mathcal{C}_\lambda^{\mathrm{Trun}}$, which was referred to as the *truncated cone with vertex* λ in [16], to be the subset of $\mathcal{C}_\lambda^{\mathrm{Norm}}$ consisting of weights μ such that the s-th entry of the atypical tuple aty_μ (cf. (2.9)) is smaller than or equal to the t-th entry of aty_μ when the atypical roots γ_s,γ_t of λ (not μ) are strongly c-related for $s<t$. Namely (recall the ρ-translated notation in (3.3))

$$\mathcal{C}_\lambda^{\mathrm{Trun}}=\{\mu\in\mathcal{C}_\lambda^{\mathrm{Norm}}\,|\,\mu^\rho_{\dot{n}_s}\leq\mu^\rho_{\dot{n}_t}\ \text{ if }\ \widehat{c}_{s,t}(\lambda)=1\ \text{ for }\ s<t\}. \tag{4.5}$$

For $\lambda=\Lambda-\sum_{s=1}^r i_s\gamma_s\in\mathcal{C}_\Lambda^{\mathrm{Norm}}$, we denote

$$|\Lambda-\lambda|=\sum_{s=1}^{r}i_s \quad \text{(called the \textit{relative level of} } \lambda). \tag{4.6}$$

As an application of Theorem 4.1 we prove the following character formula which was a conjecture put forward by van der Jeugt, Hughes, King and Thierry-Mieg in [16] as the result of in depth research carried out by the authors over several years time.

Theorem 4.2.

$$\mathrm{ch}\,V(\Lambda)=\sum_{\lambda\in\mathcal{C}_\Lambda^{\mathrm{Trun}}}(-1)^{|\Lambda-\lambda|}\chi^{\mathrm{Kac}}(\lambda). \tag{4.7}$$

Proof. First we remark that for any weight $\lambda\in P$, we have (cf. (4.2))

$$\chi^{\mathrm{Kac}}(\lambda)=0 \quad \text{if } \lambda \text{ is vanishing (i.e., not regular)}. \tag{4.8}$$

Note that for any weight λ in the truncated cone $\mathcal{C}_\Lambda^{\mathrm{Trun}}$ or in the set

$$R_\Lambda:=\{\mu\in P_+\,|\,\mu\preccurlyeq\Lambda\} \quad \text{(cf. the right-hand side of (4.3))}, \tag{4.9}$$

we have the equality of the typical tuples:

$$\mathrm{typ}_\lambda=\mathrm{typ}_\Lambda \quad \text{(cf. (3.5))}. \tag{4.10}$$

Thus λ is uniquely determined by the atypical tuple aty_λ (cf. (2.9)). Clearly for any $\mu\in\mathcal{C}_\Lambda^{\mathrm{Trun}}$, it is either vanishing (so $\chi^{\mathrm{Kac}}(\mu)=0$) or is W-conjugate under the dot action to some unique $\lambda\in R_\Lambda$, where the later fact is equivalent to the existence of a unique $\sigma\in Sym_r$ satisfying $\mathrm{aty}_{\sigma\cdot\mu}=\mathrm{aty}_\lambda$ (cf. (3.38) and (3.39)). If $\sigma(t)<\sigma(s)$ for some $1\leq s<t\leq r$. Then

$$\mu^\rho_{\dot{n}_{\sigma(t)}}=\lambda^\rho_{\dot{n}_t}>\lambda^\rho_{\dot{n}_s}=\mu^\rho_{\dot{n}_{\sigma(s)}}.$$

Thus $\widehat{c}_{\sigma(t),\sigma(s)}(\Lambda)=0$ by (4.5). Therefore $\sigma\in S^\Lambda$ by (3.40). Obviously $\mathrm{aty}_\lambda=\mathrm{aty}_{\sigma\cdot\mu}\leq\mathrm{aty}_{\sigma\cdot\Lambda}$ (cf. (3.14)), i.e., $\sigma\in S^{\Lambda,\lambda}$ by definitions (3.41) and (3.13) and the fact that

$\mathrm{aty}_{\sigma\cdot\Lambda} = \mathrm{aty}_\Lambda = \mathrm{aty}_\lambda$ (cf. (4.10)). Conversely, for any $\lambda \in R_\Lambda$ and $\sigma \in S^{\Lambda,\lambda}$, there corresponds to a unique $\mu \in \mathcal{C}_\Lambda^{\mathrm{Trun}}$ such that $\mathrm{aty}_{\sigma\cdot\mu} = \mathrm{aty}_\lambda$.

For any $\lambda \in R_\Lambda$, let $\tilde\lambda \in \mathcal{C}_\Lambda^{\mathrm{Trun}} \cap W\cdot\lambda$ be the (unique) lexical weight in the sense of Definition 3.9. Then

$$\chi^{\mathrm{Kac}}(\mu) = \chi^{\mathrm{Kac}}(\tilde\lambda) \quad \text{for} \ \ \mu \in \mathcal{C}_\Lambda^{\mathrm{Trun}} \cap W\cdot\lambda, \tag{4.11}$$

since elements of Sym_r correspond to elements of W with even parity (cf. (3.38) and (3.39)).

The above arguments have in fact shown that the right-hand side of (4.7) is equal to

$$\sum_{\lambda\in P_+:\,\lambda\preccurlyeq\Lambda} (-1)^{|\Lambda-\lambda|} \sum_{\sigma\in S^{\Lambda,\lambda}} \chi^{\mathrm{Kac}}(\tilde\lambda) = \sum_{\lambda\in P_+:\,\lambda\preccurlyeq\Lambda} (-1)^{|\Lambda-\lambda|} m(\Lambda)_\lambda \chi^{\mathrm{Kac}}(\tilde\lambda). \tag{4.12}$$

By (4.3), what remains to prove is the following: if $w(\tilde\lambda^\rho) = \lambda^\rho$ for $w \in W$, $\lambda \in R_\Lambda$, then

$$(-1)^{|\Lambda-\lambda|} = (-1)^{\ell(\Lambda,\lambda)}\epsilon(w). \tag{4.13}$$

Note that in order to obtain λ^ρ from $\tilde\lambda^\rho$ by moving all entries $\tilde\lambda^\rho_{m_s}$, $\tilde\lambda^\rho_{\dot n_s}$ for $s = 1, ..., r$ to suitable positions step by step, each time exchanging nearest neighbor entries only, the total number of movements is $N = \sum_{s=1}^r N_s$, where

$$N_s = \#([\tilde\lambda^\rho_{\dot n_s} + 1, \Lambda^\rho_{\dot n_s}] \cap \mathrm{Set}(\mathrm{typ}_\Lambda)) \quad \text{(cf. notation (3.20))}.$$

To see this, note that $\tilde\lambda$ is obtained from Λ by subtracting some atypical roots (cf. (4.4)). Thus the entries $\tilde\lambda^\rho_{m_s}$, $\tilde\lambda^\rho_{\dot n_s}$ of $\tilde\lambda^\rho$, which seat at the positions $m_s, \dot n_s$ of Λ should be moved to appropriate positions in order to make the resultant weight λ^ρ dominant. The number of steps needed for these two entries is obviously N_s.

From (3.17) and (3.16), we have $|\Lambda - \lambda| = \sum_{s=1}^r (\Lambda_{\dot n_s} - \lambda_{\dot n_s}) = \ell(\Lambda,\lambda) + N$. But $\epsilon(w) = (-1)^N$, we obtain (4.13). □

4.2. **Definitions of $\lambda_\uparrow$ and C_r.** Our purpose is to re-write (4.7) into a finite sum. Since the sum over the truncated cone $\mathcal{C}_\Lambda^{\mathrm{Trun}}$ in (4.7) is difficult to compute, we want to change this sum into several sums over some normal cones $\mathcal{C}_\mu^{\mathrm{Norm}}$ by making use of the fact that the Kac-character $\chi^{\mathrm{Kac}}(\lambda)$ is Sym_r-invariant under the dot action (cf. (3.38) and (3.39)), as a sum over a cone $\mathcal{C}_\mu^{\mathrm{Norm}}$ is easy to compute.

This will be done in two steps.

First we need to introduce more notations. Define another partial order "$\le$" on $\mathcal{C}_\Lambda^{\mathrm{Norm}}$ such that for $\lambda, \mu \in \mathcal{C}_\Lambda^{\mathrm{Norm}}$,

$$\mu \le \lambda \quad \iff \quad \text{every entry of } \mu \quad \le \quad \text{the corresponding entry of } \lambda. \tag{4.14}$$

Definition 4.3. For $\lambda \in \mathcal{C}_\Lambda^{\mathrm{Norm}}$, denote by $\lambda_\uparrow \in \mathcal{C}_\Lambda^{\mathrm{Norm}}$ the maximal lexical weight (cf. (3.28)) which is $\le \lambda$, namely,

$$\lambda_\uparrow = \max\{\mu \in \mathcal{C}_\Lambda^{\mathrm{Norm}} \mid \mu \le \lambda, \quad \text{and} \ \ \mu \text{ is lexical}\}. \tag{4.15}$$

Thus we have the equality of typical tuples: $\mathrm{typ}_{\lambda_\uparrow} = \mathrm{typ}_\lambda$ (cf. (3.5)) and the entries of atypical tuple $\mathrm{aty}_{\lambda_\uparrow}$ (cf. (3.4)) are defined by

$$(\lambda_\uparrow)^\rho_{\dot n_s} = \min\{\lambda^\rho_{\dot n_t} \,|\, s \le t \le r\} \quad \text{for} \quad s = 1, ..., r.$$

Denote by $\mathcal{C}_\lambda^{\rm Lexi}$, called the *lexical cone with vertex* λ, the subset of the truncated cone $\mathcal{C}_\lambda^{\rm Trun}$ (cf. (4.5)) consisting of lexical weights (cf. (3.28)), namely

$$\mathcal{C}_\lambda^{\rm Lexi}=\{\mu\in\mathcal{C}_\lambda^{\rm Trun}\mid \mu \text{ is lexical}\}. \tag{4.16}$$

Our first step is to change the sum over $\mathcal{C}_\Lambda^{\rm Trun}$ in (4.7) to several sums over some lexical cones $\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\Lambda)_\uparrow}$ (see Proposition 4.4).

The proof of Theorem 3.24 (cf. the arguments of proving (4.12)) and (4.8) show that (4.7) can be re-written as

$$\mathrm{ch}\,V(\Lambda)=\sum_{\sigma\in S^\Lambda}\ \sum_{\lambda\in\mathcal{C}_\Lambda^{\rm Lexi}:\,\lambda\preccurlyeq\sigma\cdot\Lambda,\,\lambda\text{ regular}}(-1)^{|\Lambda-\lambda|}\chi^{\rm Kac}(\lambda). \tag{4.17}$$

The definition of $\lambda_\uparrow$ in (4.15) shows that we have the equality of the following two sets of regular weights:

$$\{\lambda\in\mathcal{C}_\Lambda^{\rm Lexi}\mid\lambda\preccurlyeq\sigma\cdot\Lambda,\ \lambda\text{ is regular}\}=\{\lambda\in\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\Lambda)_\uparrow}\mid\lambda\text{ is regular}\}. \tag{4.18}$$

Thus (4.17) leads to the following.

Proposition 4.4.

$$\mathrm{ch}\,V(\Lambda)=\sum_{\sigma\in S^\Lambda}\ \sum_{\lambda\in\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\Lambda)_\uparrow}}(-1)^{|\Lambda-\lambda|}\chi^{\rm Kac}(\lambda). \tag{4.19}$$

The second step is to change the sum over the lexical cone $\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\Lambda)_\uparrow}$ in (4.19) to several sums over the normal cones $\mathcal{C}^{\rm Norm}_{(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow}$. This will be achieved by Lemma 4.6 below. To state lemma, we need some further notations.

Definition 4.5. Denote by C_r the subset of Sym_r consisting of permutations π which can be written as a product of *cyclic permutations* of the form

$$\pi=(1,2,...,i_1)(i_1{+}1,i_1{+}2,...,i_1{+}i_2)\cdots(i_1{+}...{+}i_{t-1}{+}1,i_1{+}...{+}i_{t-1}{+}2,...,r), \tag{4.20}$$

where $i_1,...,i_t$ are positive integers such that $\sum_{s=1}^t i_s=r$ (namely, $(i_1,...,i_t)$ is a *composition of* r). Associated to π, there is the multi-nomial coefficient

$$\binom{r}{\pi}=\frac{r!}{i_1!\cdots i_t!}. \tag{4.21}$$

4.3. **A technical lemma.** The following technical lemma is crucial in obtaining our character formula in Theorem 4.9.

Lemma 4.6. *Let* $\lambda\in\mathcal{C}_\Lambda^{\rm Lexi}$. *We have*

$$\sum_{\mu\in\mathcal{C}_\lambda^{\rm Lexi}}(-1)^{|\Lambda-\mu|}\chi^{\rm Kac}(\mu)=\frac{1}{r!}\sum_{\pi\in C_r}\binom{r}{\pi}(-1)^{\ell(\pi)}\sum_{\mu\in\mathcal{C}^{\rm Norm}_{(\pi\cdot\lambda)_\uparrow}}(-1)^{|\Lambda-\mu|}\chi^{\rm Kac}(\mu), \tag{4.22}$$

where $\ell(\pi)$ *is the length of* π *(cf. (3.43)), namely* $\ell(\pi)=\sum_{s=1}^t(i_s-1)$ *for* π *in (4.20).*

Proof. For convenience, we denote by C'_r the set of compositions of r, and denote

$$\pi_{\underline{i}}=\ \pi\text{ which is defined by (4.20) for }\ \underline{i}=(i_1,...,i_t)\in C'_r. \tag{4.23}$$

We also denote (4.21) by $\binom{r}{\underline{i}}$. For any subset S of $\mathcal{C}_\Lambda^{\rm Norm}$, denote by $S^{\rm reg}$ the regular weights of S. We define

$$\chi^{\rm sum}(S):=\sum_{\mu\in S}(-1)^{|\Lambda-\mu|}\chi^{\rm Kac}(\mu). \tag{4.24}$$

Thus

$$\chi^{\rm sum}(S)=\chi^{\rm sum}(S^{\rm reg}). \tag{4.25}$$

Note that any element $\mu\in\mathcal{C}_\Lambda^{\rm Norm}$ is uniquely determined by the atypical tuple ${\rm aty}_\mu$ (cf. (3.4)), thus there is a bijection

$$\varphi:\Lambda+\sum_{s=1}^{r}\mathbb{Z}\gamma_s\to\mathbb{Z}^r,\quad \varphi(\mu)={\rm aty}_\mu=(\mu^\rho_{\dot{n}_1},...,\mu^\rho_{\dot{n}_r}). \tag{4.26}$$

For simplicity, we use $g=\varphi(\lambda)=(g_1,...,g_r)$ to represent λ (and transfer all terminologies to g). Then the normal cone $\mathcal{C}_\lambda^{\rm Norm}$ defined in (4.4) and the lexical cone $\mathcal{C}_\lambda^{\rm Lexi}$ defined in (4.16) correspond to the following sets respectively:

$$\begin{aligned}\mathcal{C}_g^{\rm Norm}&=\{x=(x_1,...,x_r)\in\mathbb{Z}^r\,|\,x_s\le g_s,\ s=1,...,r\},\\ \mathcal{C}_g^{\rm Lexi}&=\{x\in\mathbb{Z}^r\,|\,x_1\le x_2\le\cdots\le x_s\text{ and }x_s\le g_s,\ s=1,...,r\}.\end{aligned} \tag{4.27}$$

We define the *half-lexical cone with vertex* λ:

$$\mathcal{C}_g^{\rm Half}=\{x\in\mathbb{Z}^r\,|\,x_2\le\cdots\le x_s\text{ and }x_s\le g_s,\ s=1,...,r\},$$

(i.e., we relax the condition of $x_1\le x_2$, cf. (4.16) and (4.27)). Define

$$\begin{aligned}&\mathcal{N}(g_1,...,g_r)=\chi^{\rm sum}(\mathcal{C}_g^{\rm Norm}),\quad \mathcal{L}(g_1,...,g_r)=\chi^{\rm sum}(\mathcal{C}_g^{\rm Lexi}),\\ &\mathcal{H}_{g_1}(g_2,...,g_r)=\chi^{\rm sum}(\mathcal{C}_g^{\rm Half}),\end{aligned} \tag{4.28}$$

which are the sign sums of Kac-characters over the normal cone $\mathcal{C}_g^{\rm Norm}$, the lexical cone $\mathcal{C}_g^{\rm Lexi}$ and the half-lexical cone $\mathcal{C}_g^{\rm Half}$ respectively (cf. (4.24)). Then (4.22) is equivalent to

$$\mathcal{L}(g_1,...,g_r)=\frac{1}{r!}\sum_{\underline{i}\in C'_r}\binom{r}{\underline{i}}(-1)^{\ell(\underline{i})}\mathcal{N}(g_{\underline{i}}), \tag{4.29}$$

where $g_{\underline{i}}$ is defined by

$$g_{\underline{i}}=(\ \overbrace{g_1,...,g_1}^{i_1},\ \overbrace{g_{i_1+1},...,g_{i_1+1}}^{i_2},\ ...,\ \overbrace{g_{\sum_{s=1}^{t-1}i_s+1},...,g_{\sum_{s=1}^{t-1}i_s+1}}^{i_t}\ ) \tag{4.30}$$

for $\underline{i}=(i_1,...,i_t)$. This is because from the definition (4.15), one can easily check

$$g_{\underline{i}}=\varphi((\pi_{\underline{i}}\cdot\lambda)_\uparrow)\quad\text{(cf. (4.23) and (4.26))}. \tag{4.31}$$

Denote

$$\begin{aligned}g^{(s)}&=(g_1,...,g_1,g_{s+1},...,g_r)=g|_{g_2=...=g_s=g_1}\in\mathbb{Z}^r,\\ g'^{(s)}&=(g_1,...,g_1,g_{s+1},...,g_r)\in\mathbb{Z}^{r-1},\end{aligned}$$

($g'^{(s)}$ is obtained from $g^{(s)}$ by deleting the first entry) for $s=1,...,r$. Note that

$$\mathcal{H}_{g_1}(g'^{(1)})=\sum_{s=1}^{r}\mathcal{L}(g^{(s)}). \tag{4.32}$$

To see this, observe that for any regular weight $x=(x_1,...,x_r)\in\mathcal{C}_g^{\rm Half}$ (cf. (4.25)), there is a unique $s\in[1,r]$ such that

$$x_2<...<x_s<x_1<x_{s+1}<...<x_r. \tag{4.33}$$

So x is Sym_r-conjugate (thus under the inverse mapping φ^{-1}, it is Sym_r-conjugate under the dot action, cf. (3.38), (3.39)) to

$$(x_2,...,x_s,x_1,x_{s+1},...,x_r)\in\mathcal{C}_{g^{(s)}}^{\rm Lexi}\quad\text{since }x_1<g_1.$$

Conversely every regular weight of $\mathcal{C}_{g^{(s)}}^{\rm Lexi}$ is Sym_r-conjugate under the dot action to a unique weight x of $\mathcal{C}_g^{\rm Half}$ satisfying (4.33). This proves (4.32).

By (4.32), we obtain

$$\mathcal{L}(g)=\mathcal{L}(g^{(1)})=\mathcal{H}_{g_1}(g'^{(1)})-\sum_{s=2}^{r}(-1)^{s-1}\frac{1}{s(s-1)}\mathcal{H}_{g_1}(g'^{(s)}). \tag{4.34}$$

Since the first variable x_1 in $\mathcal{C}_g^{\rm Half}$ does not relate to any other variable, when g_1 is fixed, $\mathcal{H}_{g_1}(g'^{(s)})$ is in fact $\mathcal{L}(g'^{(s)})$ with respect to the $r-1$ variables $g_2,...,g_r$. By the inductive assumption on r that (4.29) holds for $r-1$, we have

$$\mathcal{H}_{g_1}(g'^{(s)})=\frac{1}{(r-1)!}\sum_{\underline{j}\in C'_{r-1}}\binom{r-1}{\underline{j}}(-1)^{\ell(\underline{j})}\mathcal{N}(g^{(s)}_{(1,\underline{j})}),\ \ s=1,...,r, \tag{4.35}$$

where

$$(1,\underline{j})=\begin{cases}(1,j_1,j_2,...) & \text{if}\ \ s=1,\\ (j_1+1,j_2,...) & \text{otherwise,}\end{cases}\qquad \text{for}\quad \underline{j}=(j_1,j_2,...)\in C'_{r-1}.$$

Thus $(1,\underline{j})$ is a composition of r. For $\underline{i}\in C'_r$, $g^{(s)}_{\underline{i}}$ is defined by (4.30) with $g_2,...,g_s$ being set to g_1. Thus each $g^{(s)}_{(1,\underline{j})}$ has the form $g_{\underline{i}}$ for some $\underline{i}\in C'_r$, and

$$g^{(s)}_{(1,\underline{j})}\!=\!g_{\underline{i}}\Longleftrightarrow\begin{cases}i_1\!=\!1,\ \underline{j}\!=\!(i_2,i_3,...)\ \text{if}\ s\!=\!1,\ \ \text{or}\\ i_1\geq s,\ 1\!+\!j_1\!+\!...\!+\!j_s\!=\!i_1\ \text{and}\ (j_{s+1},j_{s+2},...)\!=\!(i_2,i_3,...)\ \text{otherwise.}\end{cases}$$

From this one can prove by induction on r and s that the coefficient of $\mathcal{N}(g_{\underline{i}})$ in (4.35) is $\frac{1}{i_2!\cdots i_t!}b_{s,i_1}$, where $b_{0,i_1}=0$ and

$$b_{s,i_1}=\frac{1}{i_1!}-\frac{1}{1!(i_1-1)!}+\frac{1}{2!(i_1-2)!}-...+(-1)^{i_1-s}\frac{1}{s!(i_1-s)!}\quad\text{for}\ \ 1\leq s\leq i_1.$$

Using this in (4.34), we obtain that the coefficient of $\mathcal{N}(g_{\underline{i}})$ in $\mathcal{L}(g)$ is $\frac{1}{i_1!\cdots i_t!}$. This proves (4.29) and the lemma. □

Remark 4.7. Note that a special case of (4.29) is when $g_1=...=g_r$. In this case we have

$$\mathcal{L}(g_1,...,g_1)=\frac{1}{r!}\mathcal{N}(g_1,...,g_1)\qquad\text{(cf. definition (4.28)).} \tag{4.36}$$

Remark 4.8. Using notation (4.24), formula (4.19) can be written as

$${\rm ch}\,V(\Lambda)=\sum_{\sigma\in S^\Lambda}\chi^{\rm sum}(\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\Lambda)_\uparrow}). \tag{4.37}$$

We also have

$$\chi^{\rm sum}(\mathcal{C}^{\rm Norm}_\lambda)=\sum_{\sigma\in Sym_r}\chi^{\rm sum}(\mathcal{C}^{\rm Lexi}_{(\sigma\cdot\lambda)_\uparrow})\qquad\text{for}\ \ \lambda\in\mathcal{C}^{\rm Norm}_\Lambda. \tag{4.38}$$

To prove (4.38), note that the derivation of (4.37) from formula (4.7) does not depend on how $\widehat{c}_{s,t}(\lambda)$'s are defined. Thus if we simply regard $\widehat{c}_{s,t}(\lambda)$ as zero for all s,t, then $\mathcal{C}^{\rm Trun}_\lambda$ coincides with $\mathcal{C}^{\rm Norm}_\lambda$, and S^λ becomes Sym_r. Hence (4.38) can be regarded as a special case of (4.37). Formula (4.22) can be re-written as

$$\chi(\mathcal{C}^{\rm Lexi}_\lambda)=\sum_{\pi\in C_r}\frac{1}{r!}\binom{r}{\pi}(-1)^{\ell(\pi)}\chi(\mathcal{C}^{\rm Norm}_{(\pi\cdot\lambda)_\uparrow}). \tag{4.39}$$

Thus formula (4.39) is the inverse formula of (4.38).

4.4. **Kac-Weyl type formula.** While the character formula (4.3) or (4.7) is extremely useful for understanding structural features of irreducible $\mathfrak{g}$-modules, such as their resolutions in terms of Kac-modules, it is not easy to use for purposes like determining the dimensions of irreducibles. For such purposes, a Kac-Weyl type formula is more desirable. We now derive such a formula.

For any weight $\lambda \in P$ and any subset $\Gamma \subset \Delta_1^+$, we define

$$\chi_\Gamma^{\rm BL}(\lambda) = L_0^{-1} \sum_{w \in W} \epsilon(w) e^{w(\lambda+\rho_0)} \prod_{\beta \in \Delta_1^+ \setminus \Gamma} (1 + e^{-w(\beta)}), \tag{4.40}$$

(which was referred to as the *Bernstein-Leites type character* in [16]). Here and below L_0, L_1 are defined in (2.14). For any $\lambda \in \mathcal{C}_\Lambda^{\rm Norm}$ (not necessarily regular), by (4.24) and the definition of the Kac-character $\chi^{\rm Kac}(\Lambda)$ in (4.2), we have

$$\begin{aligned} \chi^{\rm sum}(\mathcal{C}_\lambda^{\rm Norm}) &= (-1)^{|\Lambda-\lambda|} \frac{L_1}{L_0} \sum_{w \in W} \sum_{0 \le i_1, \dots, i_r < \infty} (-1)^{\sum_{s=1}^r i_s} e^{\lambda+\rho-\sum_{s=1}^r i_s \gamma_s} \\ &= (-1)^{|\Lambda-\lambda|} \frac{L_1}{L_0} \sum_{w \in W} \frac{e^{\lambda+\rho}}{\prod_{s=1}^r (1+e^{-\gamma_r})} \\ &= (-1)^{|\Lambda-\lambda|} \chi_{\Gamma_\Lambda}^{\rm BL}(\lambda), \end{aligned} \tag{4.41}$$

where $\Gamma_\Lambda = \{\gamma_1, \dots, \gamma_r\}$ defined in (2.10) is the set of atypical roots of Λ (cf. (3.2)). In deriving the last equality one has made use of the expression of L_1 in (2.14).

This together with (4.37) and (4.22) (or (4.39)) proves

$${\rm ch}\, V(\Lambda) = \sum_{\sigma \in S^\Lambda,\, \pi \in C_r} \frac{1}{r!} \binom{r}{\pi} (-1)^{|\Lambda-(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow| + \ell(\pi)} \chi_{\Gamma_\Lambda}^{\rm BL}\left((\pi \cdot (\sigma \cdot \Lambda)_\uparrow)_\uparrow\right). \tag{4.42}$$

Thus we obtain the following theorem.

Theorem 4.9. *The formal character* ${\rm ch}\, V(\Lambda)$ *of the finite dimensional irreducible* $\mathfrak{g}$-*module* $V(\Lambda)$ *is given by*

$${\rm ch}\, V(\Lambda)$$

$$= \sum_{\sigma \in S^\Lambda,\, \pi \in C_r} \frac{1}{r!} \binom{r}{\pi} (-1)^{|\Lambda-(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow| + \ell(\pi)} \frac{1}{L_0} \sum_{w \in W} \epsilon(w) w \Bigg(e^{(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow+\rho_0} \prod_{\beta \in \Delta_1^+ \setminus \Gamma_\Lambda} (1+e^{-\beta}) \Bigg). \tag{4.43}$$

where notations S^Λ, $\lambda_\uparrow$, C_r *are defined by* (*3.40*), (*4.15*) *and Definition 4.5* (*see also* (*2.10*), (*2.14*), (*4.6*), (*3.43*) *and* (*4.21*) *for other notations*).

Remark 4.10. Let $\lambda \in \Lambda + \sum_{s=1}^r \mathbb{Z}\gamma_s$. Denote the set of regular lexical weights which are $\le \lambda$ by:

$${\rm Reg}_\lambda = \{\mu \in \mathcal{C}_\Lambda^{\rm Norm} \mid \mu \le \lambda, \text{ and } \mu \text{ is regular and lexical}\}. \tag{4.44}$$

Define

$$\lambda^\uparrow = \max\{\mu \in \Lambda + \sum_{s=1}^r \mathbb{Z}\gamma_s \mid \ {\rm Reg}_\mu \ = \ {\rm Reg}_\lambda \ \text{ and } \mu \text{ is lexical}\}. \tag{4.45}$$

Thus $\lambda^\uparrow$ is obtained from $\lambda_\uparrow$ by replacing the atypical entries by (cf. (4.15))

$$(\lambda^\uparrow)^\rho_{\dot n_s} = (\lambda_\uparrow)^\rho_{\dot n_{t(s)}}, \quad \text{where } \ t(s) = \max\{t \ge s \mid \widehat{c}_{p,t}(\lambda_\uparrow) = 1 \text{ for } s \le p \le t\}, \tag*{(4.45)$'$}$$

for $s = 1, \dots, r$ (note that although $\lambda_\uparrow$ is not necessarily regular, one can still define $\widehat{c}_{p,t}(\lambda_\uparrow)$ as stated in Remark 3.8). Then in formula (4.18) (hence also in formula (4.43)), for each $\sigma \in S^\Lambda$, the $(\sigma \cdot \Lambda)_\uparrow$ can be replaced by any lexical weight η with

$(\sigma\cdot\Lambda)_\uparrow\le\eta\le(\sigma\cdot\Lambda)^\uparrow$. In particular, we have another character formula:

$$\begin{aligned}&\mathrm{ch}\,V(\Lambda)\\&=\sum_{\sigma\in S^\Lambda,\,\pi\in C_r}\frac{1}{r!}\binom{r}{\pi}(-1)^{|\Lambda-(\pi\cdot(\sigma\cdot\Lambda)^\uparrow)_\uparrow|+\ell(\pi)}\frac{1}{L_0}\sum_{w\in W}\epsilon(w)w\Big(e^{(\pi\cdot(\sigma\cdot\Lambda)^\uparrow)_\uparrow+\rho_0}\prod_{\beta\in\Delta_1^+\backslash\Gamma_\Lambda}(1+e^{-\beta})\Big).\end{aligned}\tag{4.43$'$}$$

Remark 4.11. The main difference between (4.43) and (4.43)$'$ lies in that formula (4.43)$'$ keeps the number of the distinct weights $(\pi\cdot(\sigma\cdot\Lambda)^\uparrow)_\uparrow$ to be minimal. For example when $\Lambda=0$, then $(\pi\cdot(\sigma\cdot\Lambda)^\uparrow)_\uparrow=0^\uparrow$ for all $\pi\in C_r$ and $\sigma\in S^\Lambda=\{1\}$ (See Corollary 4.12 below). But all $(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow$ for $\sigma=1$, $\pi\in C_r$ are distinct (in this case $(\sigma\cdot\Lambda)_\uparrow=0$).

Corollary 4.12. *If Λ is totally connected (cf. definition after Theorem 3.24), then*

$$\mathrm{ch}\,(\Lambda)=\frac{1}{r!L_0}\sum_{w\in W}\epsilon(w)w\Big(e^{\Lambda^\uparrow+\rho_0}\prod_{\beta\in\Delta_1^+\backslash\Gamma_\Lambda}(1+e^{-\beta})\Big)=\frac{1}{r!}\chi^{\mathrm{BL}}_{\Gamma_\Lambda}(\Lambda^\uparrow),\tag{4.46}$$

where $\Lambda^\uparrow$ is defined by the way that $\Lambda^\uparrow+\rho$ is obtained from $\Lambda+\rho$ by replacing all atypical entries (cf. (3.4)) by the largest one. In particular, by taking $\Lambda=0$ we obtain the following denominator formula

$$L_0=\frac{1}{r_0!}\sum_{w\in W}\epsilon(w)w\Big(e^{0^\uparrow+\rho_0}\prod_{\beta\in\Delta_1^+\backslash\Gamma_0}(1+e^{-\beta})\Big),\tag{4.47}$$

where $r_0=\min\{m,n\}$, $\Gamma_0=\{\gamma_1^0=\epsilon_m-\delta_1,...,\gamma_{r_0}^0=\epsilon_{m+1-r_0}-\delta_{r_0}\}$ and

$$0^\uparrow=\sum_{s=1}^{r_0}(r_0-s)\gamma_s^0=(0,...,0,1,...,r_0-1\,|\,r_0-1,...,1,0,...,0)\ \ (\textit{cf. notation (2.2)}).$$

Proof. If Λ is totally connected, then $S^\Lambda=\{1\}$ and (4.46) follows from (4.36) and (4.19) with $(\sigma\cdot\Lambda)_\uparrow$ replaced by $\Lambda^\uparrow$. Observe that when $\Lambda=0$, we have $\mathrm{ch}\,V(\Lambda)=1$ and $\Gamma_\Lambda=\Gamma_0$. Thus we obtain (4.47). □

Corollary 4.13. *If Λ is totally disconnected (cf. definition after Theorem 3.24), then*

$$\mathrm{ch}\,(\Lambda)=\frac{1}{L_0}\sum_{w\in W}\epsilon(w)w\Big(e^{\Lambda+\rho_0}\prod_{\beta\in\Delta_1^+\backslash\Gamma_\Lambda}(1+e^{-\beta})\Big)=\chi^{\mathrm{BL}}_{\Gamma_\Lambda}(\Lambda).$$

Proof. If Λ is totally disconnected, then $S^\Lambda=Sym_r$ and the result follows from the fact that (4.22) is the inverse formula of (4.38) (cf. (4.41)). □

4.5. **Dimension formula.** An important application of Theorem 4.9 is the derivation of a dimension formula.

Theorem 4.14. *The dimension $\dim V(\Lambda)$ of the finite dimensional irreducible $\mathfrak{g}$-module $V(\Lambda)$ is given by*

$$\begin{aligned}&\dim V(\Lambda)\\&=\sum_{\substack{\sigma\in S^\Lambda,\,\pi\in C_r\\ B\subset\Delta_1^+\backslash\Gamma_\Lambda}}\frac{1}{r!}\binom{r}{\pi}(-1)^{|\Lambda-(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow|+\ell(\pi)}\prod_{\alpha\in\Delta_0^+}\frac{(\alpha,\rho_0+(\pi\cdot(\sigma\cdot\Lambda)_\uparrow)_\uparrow-\sum_{\beta\in B}\beta)}{(\alpha,\rho_0)}.\end{aligned}\tag{4.48}$$

Proof. Regard an element of ε (cf. (2.12)) as a function on $\mathfrak{h}^*$ such that the evaluation of e^λ on μ is $e^\lambda(\mu) = e^{(\lambda,\mu)}$ for $\mu \in \mathfrak{h}^*$. Then $\mathrm{ch}\, V(\Lambda) \in \varepsilon$ is a function on $\mathfrak{h}^*$, and

$$\dim V(\Lambda) = \lim_{x\to 0} (\mathrm{ch}\, V(\Lambda))(x\rho_0). \tag{4.49}$$

First we calculate $\lim_{x\to 0} \chi_\Gamma^{\mathrm{BL}}(\lambda)(x\rho_0)$ (cf. (4.40)). Using

$$\prod_{\beta\in\Delta_1^+\backslash\Gamma_\Lambda} (1+e^{-\beta}) = \sum_{B\subset\Delta_1^+\backslash\Gamma_\Lambda} e^{-\sum_{\beta\in B}\beta},$$

and the well-known denominator formula

$$L_0 = \sum_{w\in W} \epsilon(w) e^{w(\rho_0)} \quad \text{(cf. (2.14))}, \tag{4.50}$$

we have

$$\begin{aligned}
\lim_{x\to 0} \chi_\Gamma^{\mathrm{BL}}(\lambda)(x\rho_0) &= \lim_{x\to 0} \sum_{B\subset\Delta_1^+\backslash\Gamma_\Lambda,\, w\in W} \epsilon(w) \frac{e^{(w(\lambda+\rho_0-\sum_{\beta\in B}\beta),x\rho_0)}}{L_0(x\rho_0)} \\
&= \lim_{x\to 0} \sum_{B\subset\Delta_1^+\backslash\Gamma_\Lambda} \frac{L_0(x(\lambda+\rho_0-\sum_{\beta\in B}\beta))}{L_0(x\rho_0)} \\
&= \lim_{x\to 0} \sum_{B\subset\Delta_1^+\backslash\Gamma_\Lambda} \prod_{\alpha\in\Delta_0^+} \frac{e^{(\alpha/2,x(\lambda+\rho_0-\sum_{\beta\in B}\beta))} - e^{(-\alpha/2,x(\lambda+\rho_0-\sum_{\beta\in B}\beta))}}{e^{(\alpha/2,x\rho_0)} - e^{(-\alpha/2,x\rho_0)}} \\
&= \sum_{B\subset\Delta_1^+\backslash\Gamma_\Lambda} \prod_{\alpha\in\Delta_0^+} \frac{(\alpha, \lambda+\rho_0-\sum_{\beta\in B}\beta)}{(\alpha,\rho_0)},
\end{aligned}$$

where the second equality follows from (4.50) and the fact that $(w(\lambda),\mu) = (\lambda, w(\mu))$ for $w\in W$, $\lambda,\mu\in\mathfrak{h}^*$. This together with (4.49) and (4.42) gives the result. □

Remark 4.15. As far as we are aware, there exists no dimension formula for the finite dimensional irreducible $\mathfrak{g}$-modules in the literature except the one for singly atypical modules given by van der Jeugt [14].

Acknowledgements. We wish to thank Professor V. Kac for suggesting important improvements to the paper. Financial support from the Australian Research Council is gratefully acknowledged. Su is also supported by NSF grant 10171064 of China, EYTP and TCTPFT grants of Ministry of Education of China.

References

[1] I.N. Bernstein, D.A. Leites, *A formula for the characters of the irreducible finite-dimensional representations of Lie superalgebras of series gl and sl*, C. R. Acad. Bulgare Sci. **33** (1980), 1049-1051.

[2] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra gl(m|n)*, J. Amer. Math. Soc. **16** (2002) 185-231.

[3] J. Brundan, *Kazhdan-Lusztig polynomials and character formulae for the Lie superalgebra q(n)*, Adv. Math. **182** (2004) 28-77.

[4] S.-J. Cheng and R.B. Zhang, *An analogue of Kostant's $\mathfrak{u}$-cohomology formula for the general linear superalgebra.* International Mathematics Research Notices **1** (2004) 31-53.

[5] J.W.B. Hughes, R.C. King and J. van der Jeugt, *On the composition factors of Kac modules for the Lie superalgebras sl(m|n)*, J. Math. Phys. **33** (1992) 470-491.

[6] V.G. Kac, *Classification of simple Lie superalgebras*, Funct. Anal. Appl. **9** (1975) 263-265.

[7] V.G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977) 8-96.

[8] V.G. Kac, *Characters of typical representations of classical Lie superalgebras*, Comm. Alg. **5** (1977) 889-897

[9] V.G. Kac, *Representations of classical Lie superalgebras*, Lect. Notes Math. **676** (1978) 597-626.

[10] I.B. Penkov and V. Serganova, *Characters of irreducible G-modules and cohomology of G/P for the Lie supergroup $G = Q(N)$*, Algebraic geometry, 7, J. Math. Sci. (New York) **84** (1997), no. 5, 1382-1412.

[11] V. Serganova, *Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra $gl(m|n)$*, Selecta Math. **2** (1996) 607-654.

[12] V. Serganova, *Characters of irreducible representations of simple Lie superalgebras*, Proceedings of the International Congress of Mathematicians 1998, Berlin, Vol. II, Documenta Mathematica, Journal der Deutschen Mathematiker-Vereinigung, pp. 583-593.

[13] J. van der Jeugt, *Character formulae for the Lie superalgebra $C(n)$*, Comm. Algebra **19** (1991) 199-222.

[14] J. van der Jeugt, *Dimension formulas for the Lie superalgebras $sl(m|n)$*, J. Math. Phys. **36** (1995) 605-611.

[15] J. van der Jeugt, J.W.B. Hughes, R.C. King and J. Thierry-Mieg, *A character formula for singly atypical modules of the Lie superalgebra $sl(m/n)$*, Comm. Alg. **19** (1991) 199-222.

[16] J. van der Jeugt, J.W.B. Hughes, R.C. King and J. Thierry-Mieg, *Character formulas for irreducible modules of the Lie superalgebras $sl(m|n)$*, J. Math. Phys. **31** (1990) 2278-2304.

[17] J. van der Jeugt and R.B. Zhang, *Characters and composition factor multiplicities for the Lie superalgebra $gl(m|n)$*, Lett. Math. Phys. **47** (1999) 49-61.

[18] Y.M. Zou, *Categories of finite-dimensional weight modules over type I classical Lie superalgebras*, J. Algebra **180** (1996) 459-482.

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia; Department of Mathematics, Shanghai Jiaotong University, Shanghai 200030, China.
E-mail address: `yucai@maths.usyd.edu.au`

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia
E-mail address: `rzhang@maths.usyd.edu.au`